

VOL. XII,
OLD SERIES.

PHILADELPHIA, OCTOBER, 1898.

VOL. I. No. 10.
NEW SERIES.

PUBLISHED IN CONJUNCTION WITH
Sajous's Annual and Analytical Cyclopædia of Practical Medicine.


PRICE OF THIS JOURNAL, SEPARATE, \$5.00 PER YEAR.

THE
Monthly Cyclopædia
OF
Practical Medicine
AND
UNIVERSAL MEDICAL JOURNAL.

EDITED BY

CHARLES E. de M. SAJOUS, M.D.,

PHILADELPHIA.



LEADING ARTICLES: "Blood Examination." "Hot-Air Treatment." "Dilatation of the Stomach." "Syphilis."

THE F. A. DAVIS CO., Publishers.

Office: 1914 and 1916 Cherry Street, Philadelphia, Pa., U. S. A.

BRANCH OFFICES: New York, U. S. A., 117 W. Forty-Second St.; Chicago, Ill., U. S. A., 9 Lakeside Building.

Exchanges and editorial matter to be sent to 2043 Walnut Street, Philadelphia, Pa.
All letters concerning subscriptions, removals, advertisements, etc., to be sent to the Publishers.
This Journal should be carefully preserved for binding as an additional volume of the Cyclopædia at the end of each year.

"Meat and bread together make the essentials of a healthful diet. Such are the facts of experience. The advancing science of later years explains them."

PROF. ATWATER.

PANOPEPTON

presents all the constituents of these two great types of food in a perfectly soluble, diffusible and absorbable form.

In sickness, the indications and necessities for a comprehensive nutrient are no less plain than in health. The expedient of pre-digesting or peptonising this food is one of the most rational, scientific and important advancements of modern medicine.

PANOPEPTON

conserves and imparts energy, sustains the system and enables it to resist the inroads of disease; quickly recruits the digestive functions to their normal power. Fairchild Bros. & Foster, New York.

PACKER'S TAR SOAP.

VEGETABLE OILS, PINE-TAR,
AND GLYCERINE.



PURE. ANTISEPTIC.
EMOLLIENT.

THE STANDARD.—Used and recommended by the **Medical Profession** for a quarter of a century in the treatment of Dermatic Diseases, for cleansing fetid discharges, for making gynecological examinations, etc.

AS A SHAMPOOING AGENT Packer's Tar Soap is undoubtedly the best known. It is extensively used, both as a delightful cleanser to preserve healthful conditions of the hair and scalp, and in treatment of dandruff and baldness.

THE PACKER MANUFACTURING CO., 81 Fulton Street. New York

HENRY'S Three Chlorides

(LIQ. FERRISENIC HENRY'S).

THE GLANDULAR STIMULANT AND
CHALYBEATE TONIC.

Tonic, stomachic, antiscorbutic,
hæmatinic and alterative.

In Anæmias, Chlorosis. A reconstructive in convalescence from any cause

Makes Quantity and Quality of Blood.

DOSE: 1 or 2 drachms.

12 oz., \$1.00.

LITERATURE ON APPLICATION.

HENRY'S TRI-IODIDES.

(LIQ. SALI-IOIDIDE HENRY'S.)

ALTERATIVE (Antirheumatic.
Antisyphilitic.

In lymphatic and visceral disturbances a uric acid solvent, and a reliable eliminant in all diathetic diseases.

8 oz., \$1.00.

1 or 2 fluidrachms in water exercises a synergistic effect over the entire range of

THE SALICYLATES AND IODIDES.

MAIZO- LITHIUM

(LIQ. LITHIUM MAIZENATE).

A POSITIVE DIURETIC.

STIMULATING, YET NON-IRRITATING.

In genito-urinary diseases, cystitis, incipient diabetes, lithæmia, urethritis, and all inflamed conditions requiring a non-irritating diuretic.

DOSE: 1 or 2 drachms

8 oz., \$1.00.

HENRY PHARMACAL CO., LOUISVILLE, KY.

BENSOLYPTUS

Schieffelin's

Alkaline Antiseptic and Prophylactic.

Bensolyptus is an agreeable alkaline solution of various highly approved antiseptics, so combined that one supplements the action of the other. Its ingredients are all of recognized value in the treatment of

Catarrhal Affections

because of their cleansing, soothing, and healing properties. Bensolyptus is highly recommended in all diseases of the nose and throat, both acute and chronic, and as a mouth-wash and dentifrice. It is also indicated for internal use in affections of the stomach and intestines where an agreeable, unirritating, and efficient antiseptic and antifermentative is required.

Pamphlets on application to

Schieffelin & Co., New York.

Schieffelin & Co.'s

(NEW YORK)

Soluble Pills and Granules.

Reliable. Soluble. Beautiful.

The Perfection of Pill Making.

Examine Them and Notice

Their Perfect Uniformity in size and weight.
The Crystal Transparency of the coating.
They are the only pills which show the precise colors of the masses.

Test Them and Prove

Their Ready Solubility.
Their Exact Conformity to their several Formulas.
The Certainty of Proper Therapeutic Effects.

Our list of Pills and Granules embraces those made according to the formulas of the U. S. Pharmacopœia; also most of those in common use among the profession.
We also manufacture pills to order.

SPECIFY Schieffelin's WHEN PRESCRIBING.

ALWAYS THE SAME. A STANDARD OF ANTISEPTIC WORTH.

Listerine.

LISTERINE is a non-toxic, non-irritating and non-escharotic antiseptic, composed of ozoniferous essences, vegetable antiseptics and benzo-boracic acid.

LISTERINE is to make and maintain surgical cleanliness in the antiseptic and prophylactic treatment and care of all parts of the human body.

LISTERINE is invaluable in obstetrics and gynecology as a general cleansing, prophylactic, or antiseptic agent.

LISTERINE is useful in the treatment of the infectious maladies which are attended by inflammation of accessible surfaces—as diphtheria, scarlet fever and pertussis.

LISTERINE diluted with water or glycerine speedily relieves certain fermentative forms of indigestion.

LISTERINE is indispensable for the preservation of the teeth, and for maintaining the mucous membrane of the mouth in a healthy condition.

LISTERINE is of accurately determined and uniform antiseptic power, and of positive originality.

LISTERINE is kept in stock by all worthy pharmacists everywhere.

Lambert's Lithiated Hydrangea.

A valuable renal alternative and antilithic agent of marked service in the treatment of Cystitis, Gout, Rheumatism, and diseases of the Uric Diathesis generally.

LITERATURE UPON APPLICATION.

LAMBERT PHARMACAL CO., ST. LOUIS.

HIGHLAND WATER



*Is the Best Tonic, Diuretic, and
Solvent Eliminator of Uric Acid:*

1. **Because** it is the softest and purest water known, containing no excess of mineral salts and no lime or organic impurities, being therefore non-irritating to the digestive organs and kidneys.
2. **Because** it contains Iron in combination with the alkalies, Sodium and Potassium in small quantities, as recommended by Professors Flint, Thomson, Draper, and others.
3. **Because** it comes from a region free from malarial germs.
4. **Because** all its mineral salts, and five of them in the exact form given in its analysis, are found in the human blood, of which they form a most essential part.

At Grocers and Druggists, or

Office, 439 FIFTH AVENUE,

NEW YORK.

Telephone, 1763-38th St.

In writing to advertisers mention Monthly Cyclopædia

SVAPNIA

OR

PURIFIED OPIUM

FOR PHYSICIANS USE ONLY.

Contains the Anodyne and Soporific Alkaloids, Codeia, Narceia and Morphia. Excludes the Poisonous and Convulsive Alkaloids, Thebaine, Narcotine and Papaverine.

SVAPNIA has been in steadily increasing use for over twenty years, and whenever used has given great satisfaction.

To PHYSICIANS OF REPUTE, not already acquainted with its merits, samples will be mailed on application.

SVAPNIA is made to conform to a uniform standard of Opium of Ten per cent. Morphia strength.

JOHN FARR, Manufacturing Chemist, New York.

C. N. CRITTENTON, Gen'l Agent, 115 Fulton St., N. Y.

To whom all orders for samples must be addressed.

SVAPNIA IS FOR SALE BY DRUGGISTS GENERALLY.

Continue to use Parke, Davis & Co.'s Anti-Diphtheritic Serum. We continue to lead in potency, efficiency and reliability of product. Remember also that P., D. & Co.'s serum is marketed in hermetically sealed bulbs (not ordinary vials and corks), which exclude the air and keep the serum strictly aseptic.

Latest literature mailed upon request.

Special Notice.—The validity of the recent patent issued by the U. S. Patent Office on diphtheria anti-toxin will be contested in the courts. Pending a decision in the case, we beg to announce that we shall, at our own expense, protect and defend physicians in any legal proceedings which may be brought as a result of the purchase of "P., D. & Co.'s" Anti-Diphtheritic Serum. Order it at any time and in any quantity without hesitation.

Parke, Davis & Co.,

Home Offices and Laboratories, Detroit, Mich.

Branches in New York, Kansas City, Baltimore, and New Orleans.

IN OVARIAN CONGESTION, HYSTERALGIA, DYSMENORRHOEA ETC.
Crush—Take with Water or Whisky.



ANTI-KAMNIA AND CODEINE TABLETS.
 (4 3/4 gr. Antikamnia, 1/2 gr. Sulph. Codeine.)
 IN LARYNGEAL NEUROSES, BRONCHIAL AFFECTIONS,
 PHTHISICAL COUGH AND GRIPPEAL CONDITIONS.
Slowly dissolve Tablet upon the tongue.

SYRUPUS ROBORANS.

Syr. Hypophos. Comp. with Quin., Mangan., and Strych.
 1-128 gr. Strychnine to teaspoonful.

Syrupus Roborans as a Tonic during Convalescence has no equal.

As a nerve stimulant and restorative in wasting and debilitating diseases, as a constructive agent in Insomnia, Pneumonia, Tuberculosis, Bronchial Asthma, Marasmus, Strumous Diseases, and General Debility, this compound has no superior.

SYRUPUS ROBORANS is in perfect solution and will keep in any climate.

Dr. T. H. STUCKY writes: In a case of Tertiary Syphilis, very anæmic, the Iodides were revolting to the stomach, being vomited when taken. Syrupus Roborans given three weeks with improvement, when the Iodide of Potassium was retained with good results.

Dr. W. O. ROBERTS says: In cases convalescing from "La Grippe" Syrupus Roborans has no equal.

For Sale by all Wholesale Druggists.

PETER'S PEPTIC ESSENCE CO.

A Powerful Digestive Fluid in Palatable Form.

Please note that Essence and Elixir Pepsin contain only Pepsin, while in Peter's Peptic Essence we have all the digestive ferments. These are preserved in solution with C. P. Glycerin in a manner retaining their full therapeutic value, which is exerted in and beyond the stomach.

It is a Stomachic Tonic, and relieves Indigestion, Flatulency, and has the remarkable property of arresting Vomiting during pregnancy. It is a remedy of great value in Gastralgia, Enteralgia, Cholera Infantum, and Intestinal derangements, especially those of an Inflammatory character. For nursing mothers and teething children it has no superior. Samples sent upon application. Express charges at your expense.

ARTHUR PETER & CO., Louisville, Ky.

EUQUININE....



Tasteless, Nontoxic QUININE

ESPECIALLY INDICATED IN CHILDREN'S CASES.

A SPECIFIC AGAINST WHOOPING-COUGH.

R. W. PHAIR & CO., Agents, 16 Platt Street, NEW YORK.

In writing to advertisers mention Monthly Cyclopædia

VEREINIGTE OHNINLABRIKEN.....


ZIMMER & CO.,

Frankfort O. M., - - Germany.

...VALIDOL

An Analeptic and Antihysteria


SPECIFIC AGAINST SEASICKNESS



THE baths to be obtained in this institution consist of the Nauheim, or Schott, bath for the various forms of Heart Disease.

Dry Superheated Air Cylinder for Rheumatism, etc., and Salt Rub Baths.

The various forms of Electricity and Massage are employed as indicated.



Rotterham House Hospital

(ESTABLISHED IN 1887),

54, 56, and 58 Isabella Street, Toronto, Canada.

A private institution for the treatment of Diseases of the Nervous System (both sexes), Abdominal Surgery, Rheumatism, and Heart Disease.

Address, HOLFORD WALKER, M.D.

WELCH'S GRAPE JUICE

CONTAINS ALL THE NUTRIENT PROPERTIES OF THE CONCORD GRAPE, WITH THE SKIN AND SEEDS REMOVED. IT IS THE BRIGHT RED BLOOD OF THE GRAPE, UNCHANGED IN THE PROCESS FROM THE LUSCIOUS CLUSTER TO YOUR PATIENT'S GLASS, AND BUT LITTLE CHANGED FROM GRAPE JUICE TO BLOOD.

WELCH'S GRAPE JUICE IS NOT A MEDICINE, IT IS NOT A STIMULANT—IT IS A LIQUID BREAD, A PERFECT FOOD. IN FEVERS AND WASTING DISEASES IT PRODUCES AN IMMEDIATE VIGOR FROM WHICH THERE IS NO REACTION.

PINT BOTTLE FREE TO ANY PHYSICIAN WHO WILL PAY EXPRESS CHARGES. 3 OZ. BOTTLE 10 CENTS. BOOKLET FREE

WELCH GRAPE JUICE CO., Westfield, N. Y.

ESTABLISHED, VINELAND, N. J., 1869.

In writing to advertisers mention Monthly Cyclopædia

ZE=AN^(N., B. & Co.)

A Concentrated Fluid Extract of
Corn Silk.

Ze-an (N., B. & Co.) is the most active diuretic of the materia medica, a solvent of uric acid and phosphatic gravel and an anodyne to the genito-urinary mucous membrane.

Ze-an (N.B.&Co.) is *par excellence* the remedy for

**NEPHRITIS
CYSTITIS**

URETHRITIS

and for **URIC ACID** and

PHOSPHATIC

GRAVEL.

Useful also for the removal of dropsical effusions.

UTEROTONIC

(N., B. & Co.)

A combination in palatable form of Squaw Vine, Black Haw, Jamaica Dogwood, Golden Seal, Black Cohosh, Cascara Sagrada, Hyoscyamus and Potassium Bromide.

UTEROTONIC (N., B. & Co.) is the most successful remedy yet offered for atonicity and irritability of the uterus and ovaries,

Dysmenorrhœa,

Menorrhagia,

Ovarian Neuralgia,

Endometritis,

and as a preparation for **Parturition,**

Increasing the force and lessening the pain of the uterine contractions.

Full Descriptive Pamphlets and Samples of the above furnished on application to

NELSON, BAKER & CO.,

Manufacturing Pharmacists,

DETROIT, MICH.



BAIRD STEAM FOMENTER

A UNIVERSAL MEDICAL
AND HOUSEHOLD APPARATUS

It consists of an alcohol heater, in connection with a boiler filled with water, to which a coil of rubber tubing is attached. When in operation, steam circulates through the coil, thus giving a hot fomentation (moist or dry) of even temperature.

The FOMENTER supplies the most efficient, economical and satisfactory method of applying both dry and moist hot fomentation ever offered the medical profession. It gives a continuous hot fomentation of even temperature. The degree of heat can be regulated to a nicety, and, if desired, the fomentation may be continued indefinitely. Patients need not be disturbed from their sleep and rest, nor exposed to sudden changes of temperature while changing the hot fomentation, as in the case of old methods.

In cases of Muscular Stiffness and Joint Diseases, the affected parts are wrapped in flannel with straight tubing wrapped around this. Fomentation may be continued at the degree of heat desired. Moist fomentation is produced by keeping the flannels moist.

The FOMENTER is also the best medium for producing medicated fumes for the treatment of all catarrhal affections of the head, throat and lungs. The medicine is placed in the water in the boiler. The warm inhalation causes the treatment to penetrate and heal the diseased parts, where the old method has no effect.

Canopies can be made of the bed blankets for use in cases of croup and diphtheria, and for vapor baths, etc. If the patient is confined to bed, steam can be conveyed under the blanket by means of a short piece of tubing. Diaphoresis can be continued as long as desired.

There should be one in every home and physicians should prescribe it to their patients. The BAIRD STEAM FOMENTER is also valuable as a sterilizer for small instruments.

Particulars furnished on application.

Price, \$1.50 to \$3.50

BAIRD BROS. & CO., No. 170 Madison St., Chicago, Ill.

In writing to advertisers mention Monthly Cyclopædia

DUOTAL (Guaiacol Carbonate)

AND CREOSOTAL

(Creosote Carbonate)

are the most eligible and effective of germicides, and the most vigorous neutralizers of microbic poisons in the intestinal tract. The remarkable results obtained in tuberculosis, typhoid fever, rheumatoid arthritis, etc., are explainable by the great powers of elimination of toxalbumins that these remedies have been proven to possess. They are themselves neutral, non-toxic, and absolutely free from all caustic and irritant qualities.

At the Clinic of Professor E. VON LEYDEN, at Berlin, Creosotal has been used in a large number of cases of *phthisis pulmonum* with very good results. It agreed well with all the patients. It had a very favorable effect upon the night sweats and the fevers; and the cough and expectoration either disappeared entirely, or were markedly improved. The body weight increased in most cases.

Dr. PAUL JACOB, Physician-in-Chief of Professor Von Leyden's Clinic, whose report is published in full in the *Charité Annalen*, Berlin, 1897, claims that **Creosotal has a specific action in phthisis, and that every case that is not too far advanced can be greatly benefited by its use.**

ORPHOL (Betanaphthol-Bismuth-

Von Heyden)

The employment of Orphol, which is a neutral, odorless, tasteless and non-toxic powder, is indicated in all fermentative gastro-intestinal processes, in ptomaine poisonings, gastro-enteric catarrhs, typhoid fever, etc. **Practical Intestinal Antisepsis** can be effected and maintained by its use. Unlike opium, tannin, etc., Orphol in no way interferes with the digestion, so that patients suffering from dyspepsia bear it well.

Betanaphthol-Bismuth is antiseptic as well as astringent, and is therefore far superior to the new tannin compounds recently introduced; for tannin is well known not to possess any bactericide action at all. Four or five 15-grain doses of Orphol will usually cure the very worst cases of diarrhoea; and in cholera infantum 2 to 5 grains administered every three or four hours act admirably.

Orphol is soothing to the irritated and inflamed intestinal mucous membrane, besides acting as a continuous disinfectant. It does away with the dangers of caustic or poisonous antiseptic substances, such as carbolic acid, naphthol, resorcin, the bichloride of mercury, etc., and the use of complicated and uncertain diarrhoea and cholera mixtures.

XEROFORM (Tribromphenol-

Bismuth-

Von Heyden)

Xeroform is the ideal substitute for iodoform, being an antiseptic, desiccating and deodorizing agent which is odorless, non-poisonous and non-irritating, with very powerful antibacterial properties. Professors Hueppe and Fasano, Dr. Reynders, and others, have found it to be a very effectual intestinal antiseptic in cholera, intestinal tuberculosis, etc.; but its chief application is as an external antiseptic in the place of iodoform. Xeroform has been employed by Drs. Heuss, Cumston, Grünfeld, Beyer, Metall, Paschkis, and many other observers, in the most varied surgical affections; for operative procedures, amputations, enucleations, and cancer operations; for suppurative bursitis, lymphadenitis, and alveolar periostitis; for paronychias, deep abscesses, and suppurating buboes. In venereal diseases it has been extensively employed in the treatment of chancroid; and in dermatology for impetigo, furunculosis, sycosis, eczema and pruritus. It has also been successfully used in ophthalmic and gynaecological practice, and for insufflation into the nose and ear. Its fine pulverization enables the physician to cover a large surface with a very small quantity of the drug.

Literature furnished on application. **SCHERING & GLATZ, 58 Maiden Lane, New York,**
Sole Agents for the United States.

THE VIRGINIA MEDICAL SEMI-MONTHLY.

125

Editorial.
(Richmond)
Medical Journal.
Medical Review.

[May 27]

Editorial.

of Meeting.
Having arrangements.
Sanitary Medicine
(Chicago, Toledo).
Register.
Medicine and Veterinary

Maltzyme Preparations.

Introduced by the Malt-Diastase Co., of New York city, are unquestionably of the very best of the diastatic preparations on the market. Although they take the lead from the laboratory standpoint, as shown by reports from Dr. Willis G. Tucker, Professor of Chemistry, Albany Medical College, Dr. Geo. C. Diekman, Ph. G., Professor of Theoretical and Practical Pharmacy, New York College of Pharmacy, and other laboratory workers of like ability, they are receiving the more valuable approval of clinical endorsement. This enzyme has a direct value as a nutrient; and its diastatic property is marked in bringing the starch in food into a soluble condition, and converting it into easily assimilable forms of dextrin and sugar. A striking quality of one of the most valuable preparations—Maltzyme with Cod Liver oil—consists in its fluidity, the readiness with which it is poured from the bottle and its miscibility with water. Try it in suitable cases; and report the results.

Beer as a Remedy

Typhoid Fever

During convalescence from fever, when the heart and stomach have been weakened and the digestion is weak, malt liquors properly brewed increase the appetite, digestion and nutrition. (VARIOUS TEXT BOOKS ON THERAPEUTICS.)

American Physicians have at their disposal the best beer in existence

Tannhæuser and Old Stock Lager

THE WORLD'S JUDGMENT OF OUR PRODUCT

Philadelphia, 1876, Two Medals; Paris, 1878, Grand Prize; Brussels, 1888, Grand Gold Medal and Diploma of Honor; Paris, 1889, Grand Prize; Chicago, 1893, Four Prizes; Antwerp, 1894, Grand Prize; Amsterdam, 1895, Grand Prize; Brussels, 1897, Grand Prize.

The Bergner & Engel Brewing Co., Philadelphia

In writing to advertisers mention Monthly Cyclopædia

Phenalgin

**NON TOXIC, HYPNOTIC,
Analgesic, Antipyretic.
NON DEPRESSANT, STIMULANT.**

FORMULA

Ammonio-Phenylacetamide, pure white.
Made from C. P. material.
A stable and uniform product.

R Phenalgin (Etna), gr. 10.
Make one powder for
painful menstruation.

R Phenalgin (Etna), gr. 5.
Make one powder for
insomnia.

**Generous Samples of
Powder or Tablets of
Pure Phenalgin Mailed
Free to Physicians.**

ETNA CHEMICAL CO.

313 WEST STREET,

NEW YORK, U.S.A.

S GLASS & METAL S
HIGHEST GRADE
I G FOR PHYSICIANS, I G
DENTISTS ETC. N S
J.W. RUSSELL & CO. N S
MFR'S 294 PEARL ST, NEW YORK.
WRITE FOR PRICE LIST.

Send for the interesting and beautiful book entitled
**FACTS AND SUGGESTIONS REGARDING THE PI-
NELLAS PENINSULA, FLORIDA.** It is Illustrated
with 75 Beautiful Half-Tone Engravings, nearly all
of them from Photographs. Handsomely Printed on
Fine Enameled Paper. Crown Octavo. 132 Pages.
Price, 25 cents, postage prepaid. **THE F. A. DAVIS**
Co., Publishers, 1914-16 Cherry St., Phila., Pa.

AN ANTISEPTIC DRESSING IN AN ANTISEPTIC PACKAGE.

PROGRESS IN SURGERY

has been largely due to antiseptic preparations and aseptic methods. One great want of the surgeon and general practitioner has been a surgical dressing that was thoroughly antiseptic, put up in a thoroughly antiseptic container and applicable wherever inflammation was present. Unguentine in the collapsible tube fills this want. Observing physicians will admit that there is danger of contaminating an antiseptic ointment when put up in jars or other containers where a large surface is exposed when the lid or cover is left off or not securely replaced. Acting upon the suggestions of prominent physicians we take pleasure in offering to the profession Unguentine in the collapsible tube, which gives an antiseptic dressing put up in a thoroughly antiseptic container.

Its advantages are many. The physician can carry it in his case without danger of soiling either instruments, labels, or any of its contents. It is most convenient, as the ointment can be easily applied on skin or bandage without the use of a spatula or other instrument; and on account of size of the package you can always have the best surgical dressing for minor work with you.

It is the most economical, as it is the least expensive.

The tube holds about two ozs. and costs 25 cents. From the numerous letters of commendation we have already received from the profession we feel sure that the friends and users of Unguentine are pleased with this new container.


To introduce Unguentine to you in the collapsible tube we will be pleased to send you, upon request, one tube, free of expense.

THE NORWICH PHARMACAL COMPANY,

SOLE MANUFACTURERS,

NORWICH, NEW YORK.

In writing to advertisers mention Monthly Cyclopædia



Maltine

MALTINE is not merely "malt," nor is it a mere "extract of malt," nor an "essence of malt."

MALTINE is the most highly concentrated extraction of all the nutritive and digestive properties of Wheat, Oats, and Malting Barley.

It has stood alone and unrivalled throughout the world in the therapeutic field for more than twenty years, despite the most strenuous efforts of the ablest pharmaceutical manufacturers to produce a preparation approaching it in medicinal value, elegance, palatability, and stability.

"Malt" is not "MALTINE."

"Extract of Malt" is not "MALTINE."

"Essence of Malt" is not "MALTINE."

"MALTINE" must be designated to get "MALTINE."

DOCTOR Do You Cure Your Goitre Patients?

I cure all of mine and receive from \$25 to \$50 from each of them. The cost of treating a case will not exceed \$1.00. Enclose a 2-cent stamp for particulars.

Lock Box 604.

F. E. MAY, M. D., Bloomington, Ill.

JOHN WELSH DULLES
EDWARD W. CRITTENDEN
DAVID SWOPE

Telephone 5149

DULLES & CRITTENDEN,

INSURANCE,

NO. 312 WALNUT STREET,
PHILADELPHIA.

FIRE
LIABILITY
BOILER
PLATE GLASS



CAPSULES.

		PER DOZ.	
No. 53A	{ 12 in Box . . .	\$2.25	
" 53	{ 10 Minims { 24 " . . .	4.25	
" 54	{ Size. { 36 " . . .	6.25	
" 54C	{ 100 " . . .	15.00	

"PERLOIDS,"

OR IMPROVED FRENCH "PERLES."

Cheaper and Better than the Imported.

		PER DOZ.	
No. 421A	{ 5 Minims { 40 in Vial . . .	\$4.75	
" 421B	{ Size. { 80 " . . .	9.00	
" 421C	{ 100 " . . .	10.20	

A trial bottle or dozen sent prepaid on receipt of price.

H. PLANTEN & SON (Established 1836), New York.

Manufacturers of over 400 varieties of
FILLED AND EMPTY CAPSULES.

In writing to advertisers mention Monthly Cyclopædia

**Uniformly Effective, Agreeable and Lasting,—the
Standard Preparation of Erythroxylon Coca**

During past 30 years
most popularly used
Tonic-Stimulant in
Hospitals, Public and
Religious Institutions
everywhere.



We have received
over 7000 written
endorsements from
PROMINENT PHYSI-
CIANS in Europe and
America.

"MARIANI WINE"

FORMULA : The concentrated extract—the aromatic principle of the fresh Coca Leaf,
blended with a special quality of grape juice of southern France.

DOSE : Wine-glassful three times a day, or more or less at Physician's discretion.

Nourishes - Fortifies - Refreshes
AIDS DIGESTION - STRENGTHENS THE SYSTEM

AGREEABLE TONIC-STIMULANT WITHOUT UNPLEASANT REACTION.

To avoid disappointment please specify "Vin Mariani."

SOLD AT ALL PHARMACIES.

PARIS: 41 Boulevard Haussmann.
LONDON: 239 Oxford Street.
MONTREAL: 28 Hospital Street.

MARIANI & CO., 52 W. 15th St., New York

AXION ELASTIC TRUSS.

WORN NIGHT AND DAY WITH COMFORT. HOLDS THE
WORST RUPTURE WITH EASE.



We are now manufacturing a superior grade Silk Elastic Webbing. This
Webbing, made of selected rubber, wound with Silk, has a strength, durability, and
softness of texture unequalled. This Webbing is used only in our **AXION ELASTIC TRUSS.**

Our **New Axion Pad** has now been in successful operation for six months, and is giving universal
satisfaction. It is the most durable and comfortable soft pad made. It can be made **firm** or **soft** at any
time by simply turning a screw. Perfectly adjustable in every way—it is adapted to every variety of
Hernia.

The **AXION ELASTIC TRUSS** holds rupture perfectly, exercising and resting, and is worn
with the greatest comfort by the patient.

SEND FOR ILLUSTRATED CATALOGUE.

G. V. HOUSE MFG. CO.,

744 Broadway, New York City.

Send for our New 28-Page Descriptive Catalogue. Mailed to any
address promptly on request. **THE F. A. DAVIS COMPANY,**
Medical Publishers, 1914-16 Cherry Street, Philadelphia, Pa.

The American
THERAPIST

A Monthly Record of Modern
Therapeutics.

Original, Practical,
Independent, Interesting.

SUBSCRIPTIONS SOLICITED.

THE AMERICAN THERAPIST PUBLISHING CO.,
Lock Box 1170, New York.

\$1.00
PER YEAR

Sample Copies, 10 cts.

In writing to advertisers mention Monthly Cyclopædia

CHIONIA

THE HEPATIC STIMULANT

INDICATED IN

Diseases Caused by Hepatic Torpor.

Does not purge, per se, but under its use the Liver and Bowels gradually resume their normal functions.

DOSE—ONE TO TWO FLUID DRACHMS, THREE TIMES A DAY.

PEACOCK'S BROMIDES

THE STANDARD SEDATIVE

INDICATED IN

Congestive, Convulsive and Reflex Neuroses.

Absolutely uniform in purity and therapeutic power, produces clinical results which can not be obtained from the use of commercial bromide substitutes.

DOSE—ONE TO TWO FLUID DRACHMS IN WATER, THREE TIMES PER DAY.

PEACOCK CHEMICAL COMPANY, St. Louis, Mo.

—AND—

36 BASINGHALL ST., LONDON, ENGLAND.

SENG

FOR

**INDIGESTION, MALNUTRITION, PHTHISIS,
AND ALL WASTING DISEASES.**

DOSE—One or more teaspoonfuls three times a day. For babies, ten to fifteen drops during each feeding.

CACTINA PILLETS

FOR **ABNORMAL HEART ACTION.**

DOSE—One Pillet every hour, or less often as indicated.

SULTAN DRUG CO., St. Louis and London.

In writing to advertisers mention Monthly Cyclopædia

THE GREAT FACT IN MODERN MEDICINE:

"The Blood is the Life,"

And Where Nature fails to make Good Blood,

WE CAN INTRODUCE IT.

BOVININE is Bovine Blood Unaltered from the Arteries of the Bullock;
The Universal Auxiliary of Modern Medicine and Surgery,
and the TRUE "ANTITOXIN" of Healthy Nature.

In the more enlightened progress of Modern Medicine, "Blood-letting" has given place to Blood-*getting*.

Aye! Get Good Blood—but How? Not by the Alimentary Process. It has already failed to do its work (else the patient would not be sick); and in acute disease must not even be allowed to do the work it can. Stimulate as you will, the whole sum of the patient's alimentary power when fully forced into play, is unable to keep up the nourishing and supporting contents of the blood. There is absolutely but one thing to do; and, thank God, that can be done, usually with success, as ten-thousand-fold experience has proved. That one thing is this: where Nature fails to PRODUCE good and sufficient Blood, WE CAN INTRODUCE IT from the arteries of the sturdy bullock, by the medium of BOVININE.

The vital activity of this living blood conserve rests on no man's assertion: it speaks for itself, to every properly equipped physician who will test its properties microscopically, physically, or therapeutically.

TRY IT IN PRACTICE.

TRY it in Anæmia, measuring the increase of red cells and hæmaglobin in the blood as you proceed, together with the improving strength and functions of your patient.

Try it in Consumption, with the same tests from week to week.

Try it in Dyspepsia or Malnutrition of young or old, and watch the recuperation of the paralysed alimentary powers.

Try it in Intestinal or gastric irritation, inflammation, or ulceration, that inhibits food itself, and witness the nourishing, supporting and healing work done entirely by absorption, without the slightest functional labor or irritation; even in the most delicate and critical conditions, such as Typhoid Fever and other dangerous gastro-intestinal diseases, Cholera Infantum, Marasmus, Diarrhœa, Dysentery, etc.

Try it per rectum, when the stomach is entirely unavailable or inadequate.

Try it by subcutaneous injection, when collapse calls for instantaneous blood supply—so much better than blood-dilution!

Try it on Chronic Ulceration, in connection with your antiseptic and stimulating treatment (which affords no nourishment) and prove the certainty and power of topical blood nutrition, abolishing pus, stench, and PAIN, and healing with magical rapidity and *finality*.

Try it in Chronic Catarrhal Diseases; spraying it on the diseased surfaces, with immediate addition of peroxide of hydrogen; wash off instantly the decomposed exudation, scabs and dead tissue with antiseptic solution (Thiersch's); and then see how the mucous membrane stripped open and clean, will absorb nutrition, vitality and health from intermediate applications of pure bovine.

Try it on the Diphtheritic Membrane itself, by the same process; so keeping the parts clean and unobstructed, washing away the poison, and meanwhile sustaining the strength independently of the impaired alimentary process and of exhaustive stimulants.

Try it on anything, except plethora or unreduced inflammation; but first take time to regulate the secretions and functions.

Try it on the patient tentatively at first, to see how much and how often, and in what medium, it will prove most acceptable—in water, milk, coffee, wine, grape, lemon or lime juice, broth, etc. A few cases may even have to begin by drops in crushed ice.

A New Hand-book of Hæmathery for 1898, epitomizing the clinical experience of the previous three or four years, from the extensive reports of Hospital and private practice. To be obtained of


THE BOVININE COMPANY, 75 W. Houston Street, New York.

LEEMING, MILES & Co., Montreal, Sole Agents for the Dominion of Canada.

In writing to advertisers mention Monthly Cyclopædia

You Can Boil Our Aseptic Syringe

AND KEEP IT CLEAN WITH THE LEAST AMOUNT OF EFFORT.



IT HAS { No Caps,
No Washers,
No Receptacles for Dirt.

IT IS { Handsome,
Durable,
Easily Adjusted.

IT DOES { Not Leak,
Not Dry Out,
Not Get Out of Order.

Together with the Smallest,
IT COSTS Handsomest Aluminum Case,
\$3.00 that holds 6 Tubes of our Soluble
Hypodermic Tablets and Two
Needles.

SHARP & DOHME,

CHICAGO.


BALTIMORE.

NEW YORK.

*The Best Medical
Electrical Apparatus
In the World.*

**TREATISE ON ELECTRO-THERAPY
FREE.**

300 pages. 450 Illustrations.



Giving complete electrical technique of such operations as Reduction of Strictures; Treatment of Dysmenorrhœa; Removal of Superfluous Hair, Warts, Moles, Nævi, &c.; Cataphoresis; Electro-Diagnosis; Metallic Electrolysis; also giving prices and descriptions. We are desirous of placing this *educator* in the hands of every physician in the land, and will send it **post-paid, without charge**, if you will write and ask for it.

McINTOSH BATTERY AND OPTICAL CO.,

521-531 Wabash Avenue, CHICAGO, ILL.

In writing to advertisers mention Monthly Cyclopædia

THE BEST PREPARED FOOD.

FOR THE BABY, THE INVALID, THE CONVALESCENT, THE AGED.

HIGHEST AWARDS WHEREVER EXHIBITED.

IT CONTAINS NO CANE SUGAR, NO GLUCOSE, NO MALT.

IMPERIAL GRANUM

NOURISHING STRENGTHENING.

PURE PALATABLE SAFE RELIABLE.

IS THE STANDARD PREPARED FOOD

NEARLY ALWAYS SUCCESSFUL WHEN ALL OTHER KINDS OF NOURISHMENT HAVE FAILED.

SAMPLES FOR CLINICAL TEST SUPPLIED TO PHYSICIANS AND TRAINED NURSES ON REQUEST.

OF TO-DAY AND WILL CONTINUE TO YIELD SATISFACTORY RESULTS IN NUTRITION FAR INTO THE FUTURE, BECAUSE ITS MERITS HAVE BEEN PROVED BY CLINICAL SUCCESS IN THE PAST.

Shipping Depot, JOHN CARLE & SONS, 153 Water Street, New York.
SOLD BY DRUGGISTS EVERYWHERE.

IMPERIAL GRANUM PREPARED FOOD FOR INVALIDS AND CHILDREN

Lycetol

The Uric Acid Solvent and Anti-arthritis.

An Important Advance in Gonorrhoea Therapeutics.

Protargol

Iodothyrene

The Active Principle of the Thyroid Gland.

The Modern Hypnotic.

Trional

Salophen

The Safest, Most Agreeable and Reliable Anti-rheumatic.

Send for samples and literature to

FARBENFABRIKEN OF ELBERFELD CO., 40 Stone St., New York.

Selling agents for the Bayer Pharmaceutical Products:

Aristol, Creosote Carbonate (Creosotal), Euphrophen, Ferro-Somatose, Guaiacol Carbonate (Duotal), Hemicanrin, Heroin, Iodothyrene, Lacto-Somatose, Losophan, Lycetol, Phenacetin, Piperazine-Bayer, Protargol, Quinalgen, Salicylic Acid, Salophen, Somatose, Sulfonal, Tannigen, Tannopine, Trional.

In writing to advertisers mention Monthly Cyclopædia

The only prominent Emulsion of Cod-Liver Oil introduced directly to the medical profession.
It is advertised exclusively in medical journals.

HYDROLEINE.

(HYDRATED OIL.)

Produces rapid increase in Flesh and Strength.

FORMULA.—Each Dose contains			Recommended and Prescribed by EMINENT PHYSICIANS Everywhere. It is pleasant to the Taste and acceptable to the most delicate Stomach.
Pure Cod Liver Oil.....80 m. (drops)	Soda		
Distilled Water.....35 "	Salicylic Acid ..1-4 "		
Soluble Pancreatin 5 Grains.			

IT IS ECONOMICAL IN USE AND CERTAIN IN RESULTS.

HYDROLEINE (Hydrated Oil) is not a simple alkaline emulsion of oleum morrhuae, but a hydro-pancreated preparation, containing acids and a small percentage of soda. Pancreatin is the digestive principle of fatty foods, and in the soluble form here used, readily converts the oleaginous material into assimilable matter, a change so necessary to the reparative process in all wasting diseases.

Lautenbach's Researches on the functions of the liver would show the beautiful adjustment of therapeutics in preparation of Hydroleine, furnishing, as it does, the acid and soda necessary to prevent self-poisoning by re-absorption of morbid tubercular detritus and purulent matters into the general circulation.

In Wasting Diseases the most prominent symptom is *emaciation*, the result of starvation of the fatty tissues of the body as well as the brain and nerves. This tendency to emaciation and loss of weight is arrested by the regular use of *Hydroleine*, which may be discontinued when the usual average weight has been permanently regained.

The following are some of the diseases in which **HYDROLEINE** is indicated.

Phthisis, Tuberculosis, Bronchitis, Catarrh, Cough,
Scrofula, Chlorosis, General Debility, etc.

To Brain Workers of all classes, *Hydroleine* is invaluable, supplying as it does, the true brain-food, and being more easily assimilated by the digestive organs than any other emulsion.

The principles upon which this discovery is based have been described in a treatise on "The Digestion and Assimilation of Fats in the Human Body," by H. C. BARTLETT, Ph. D., F. C. S., and the experiments which were made, together with cases illustrating the effect of Hydrated Oil in practice, are concisely stated in a treatise on "Consumption and Wasting Diseases," by G. OVEREND DREWRY, M. D.

COPIES OF THESE WORKS SENT FREE ON APPLICATION.

— Sold by all Druggists at \$1.00 per Bottle. —

C. N. CRITTENTON,

SOLE AGENT FOR THE UNITED STATES.

115 FULTON STREET, N. Y.

A Sample of Hydroleine will be sent free upon application, to any physician (enclosing business card) in the U. S.

Please mention this Journal.

In writing to advertisers mention Monthly Cyclopædia

THE MONTHLY CYCLOPÆDIA OF PRACTICAL MEDICINE.

Vol. XII.
Old Series.

PHILADELPHIA, OCTOBER, 1898.

Vol. I. No. 10.
New Series.

TABLE OF CONTENTS.

PAGE	PAGE	PAGE
BLOOD EXAMINATION. Editorial Journal of the American Medical Association; John Lovett Morse, F. P. Henry, Limbeck, G. D. Head, Thomas Turnbull (Jr.), Hans Kohn, G. Lovell Gul-land..... 361	MORPHINE POISONING. C. W. Williams..... 394 Treatment..... 394	SYPHILIS. 375 Symptoms. D. W. Montgomery, C. W. Hitchcock, Rubino, Legrain, Eugene Fuller, J. A. Fordyce, W. Morres, M. Dieulafoy, Sidney Phillips, Kirk, W. W. Whitaker, v. Duhring..... 375
DIARRHŒA, CHRONIC. Allen A. Jones..... 390 Etiology..... 390 Treatment..... 390	PROSTATITIS, CHRONIC CATARRHAL. H. M. Christian..... 394 Diagnosis..... 394	Diagnosis. C. F. Marshall, E. G. Janeway..... 378
DYSMENORRŒA. G. Granville Bantock..... 391 Etiology..... 391 Treatment..... 391	PROTARGOL, Editorial Edinburgh Medical Journal..... 395 Treatment..... 395	Etiology. Winkler, Niessen, J. G. Adami, M. Delore..... 379
EUCAINE. Lilienthal..... 392	PULMONARY TUBERCULOSIS. Tamplough..... 395 Treatment..... 395	Prognosis. Krauss, Campana..... 380
GLANDULAR FEVER IN CHILDREN. Pfeiffer..... 392 Symptoms..... 392	RECTAL PROLAPSE. O. Helms..... 396 Treatment..... 396	Treatment. Troisfontaines, Pediatrics, Manassein, Buret, Cartier, E. Lang, Dabney, Lukasiewicz, Whitla, C. F. Marshall, Jullien, Gilbert and Fournier, Justus, Fournier, Colombini and Gerulli, Cooper, Lyman Ware, Fruitnight, Ramón Gutiérrez, Keyes..... 397
GOLF. Irving C. Rosse..... 393	SEROUS PLEURISY. Proserowsky..... 396 Treatment..... 396	TYPHOID FEVER. Henry W. Bettman..... 397 Treatment..... 397
HOT-AIR TREATMENT. C. H. Frazier, E. L. Morse, A. Graham Reed, S. Solis-Cohen, J. T. Rugb, Alice M. Seabrooke..... 396	SOMATOSE. Joachim..... 396	TYPHO-MALARIAL FEVER. A. Crombie..... 397
HYPERTRICHOSIS. Eduard Schiff, Leopold Freund..... 393 Treatment..... 393	SORE NIPPLES, THE PREVENTION OF. J. Milton Mabbott..... 396	UTERINE CARCINOMA. J. H. Etheridge..... 398 Treatment..... 398
LACTOPHENIN AS HYPNOTIC. Christiansi..... 394	STARCH, SALIVARY DIGESTION OF. W. G. Aitchison Robertson..... 397	VARICOSE VEINS. Cumstom..... 398
	STOMACH, DILATION OF 370 Symptoms. W. H. Broadbent, S. B. Fowler, Fenger..... 370 Diagnosis. Pepper, Stengle..... 372 Etiology. Pepper, Stengle..... 372 Treatment. J. A. Storck, Lyman, Broadbent, Boardman Reed..... 373	NEW BOOKS RECEIVED. 399 MONOGRAPHS RECEIVED. 399 EDITORIAL STAFF 400

Cyclopædia of the Year's Literature.

BLOOD EXAMINATION.

In qualitative analysis,¹ the amount of hæmoglobin, estimated upon a percentage-basis, is the true index to the richness of the blood. Other ingredi-

ents, notably albumin, may aid somewhat; but, while the latter may be increased in the blood of fairly-healthy

¹ Jour. Amer. Med. Assoc., August 13, '98.

subjects' with considerable rapidity, hæmoglobin-increase is comparatively slow, and for that reason is a better indicator, regardless of the time of examination. The count of the number of red corpuscles, aside from the specific diseases of the blood itself, cannot be regarded as of such value as formerly; for instance, in neurasthenic states, it has lately been demonstrated that counts of 6,000,000 or 7,000,000 per cubic centimetre are not infrequent, yet the physical condition of the patient, together with his hæmoglobin, is below par. The determination of hæmoglobin is of little value from a prognostic standpoint in the various blood-affections proper; but in the simple anæmias, from whatever cause, it possesses decided importance.

In very few diseases, if any, does anæmia occur with such rapidity as in malarial fevers. With the exhibition of quinine, the majority of cases (the intermittents at any rate) become, not cured, but convalescent, and the problem now changes to that of improving the nutrition; in other words, the abolition of an anæmia necessarily present. The usual tonic treatment—iron, quinine, strychnine, Fowler's solution—is instituted, and it becomes necessary to note the effect, if any, of our drugs, especially iron, that the dosage may be regulated and the patient not given more than can be utilized. This can be best done and in a truly scientific manner by determining the hæmoglobin percentage. It need not be done oftener than once a week, and even a slight increase during that time will show to the physician that his patient is progressing in the right direction.

In syphilis there is always, in the active stage of the disease, more or less anæmia. If, to a syphilitic who has not

been treated with inunctions or injections, such treatment is instituted, a rapid fall in the percentage of hæmoglobin usually takes place, often as high as 10 to 20 per cent. Under active treatment, if not pushed to a great degree, the hæmoglobin will return to its original percentage, and under a careful administration of mercurials even increase; so that by estimating the hæmoglobin at frequent intervals, and changing the dose to suit the condition of the hæmoglobin, one may soon be able to regulate the dosage as to bring about continuous improvement in the patient. This peculiar behavior of the hæmoglobin to mercurials does not occur except in syphilis, and in such a case is a valuable diagnostic as well as prognostic point.

John Lovett Morse² asserts that the blood of infants under 2 years of age differs in certain of its characteristics from that of adults. The number of red corpuscles is about the same or a little larger, averaging a little over 5,000,000 per cubic millimetre. The number of white corpuscles per cubic millimetre is somewhat larger, averaging from 10,000 to 12,000. The relative proportions of the various forms of leucocytes are also considerably different. The proportion of mononuclear, or unripe forms, is about three times as great as in adult life, while that of the polynuclear neutrophiles, or overripe form, is only half as large. The mononuclear cells, moreover, are not merely lymphocytes, but vary much not only in the size of the cell as a whole, but also in the size of the nucleus and in the amount of protoplasm. Finally, an increase in the number of eosinophilic cells, even if con-

² Boston Med. and Surg. Jour., April 22, '97.

siderable, seems to be of less significance.

Anæmia of various sorts is very common in children. This is because the tissue-changes in them are more rapid, as the old tissues have not only to be nourished and replaced, but new ones formed. Hence any disturbance of nutrition is more serious and results in more rapid and significant changes in the blood.

Leucocytosis develops more quickly and to a higher degree in children than in adults. While in adults the increase of white cells is almost entirely in the polynuclear neutrophiles, this is not the case in children. In them the leucocytosis is sometimes due to the increase of lymphocytes, sometimes to that of the large mononuclear forms, sometimes to that of the polynuclear neutrophiles, and sometimes even to that of the eosinophiles. According to Weiss, the lymphocytes are much increased proportionately in affections of the gastro-enteric tract, while the mononuclear cells of all sorts, as well as the transition forms, are proportionately increased in those of the respiratory tract.

The writer has examined the blood in twenty cases of active, uncomplicated rickets in infants under 2 years of age. The blood was in every case taken from the lobe of the ear, and examined with a Thoma-Zeiss apparatus. It was diluted (1 to 200) with a 3-per-cent. salt solution colored with methylene-blue. The red corpuscles in $\frac{1}{10}$ cubic millimetre and the white corpuscles in 2 cubic millimetres were counted. The hæmoglobin was estimated with a von Fleischl instrument. Coverslips were in all cases made at the same time, hardened in equal parts of alcohol and ether, and stained with Ehrlich's "triple stain." A differential count of at least 500 white

corpuscles was then made, the classification recommended by Ehrlich being used. As far as possible, the blood was taken about noon, and in most cases about three hours after food. The results obtained in these cases, together with those obtained by others, seem to justify the following conclusions: Most cases of rickets are accompanied by anæmia. This anæmia may be of any form and of any grade of severity. The severity of the anæmia varies in a general way with the severity of the process. The most common form is that in which the number of red corpuscles is normal or nearly normal, and the percentage of hæmoglobin both absolutely and relatively diminished. The anæmia may or may not be accompanied by leucocytosis. Leucocytosis occurs more frequently in the cases with splenic tumor than in those without. It may be due to an increase in any or all of the varieties of white corpuscles. The specific gravity varies with the amount of hæmoglobin. Finally, there is no form of anæmia found in rickets which may not be found in other conditions, and no form of anæmia found in other conditions which may not be found in rickets.

F. P. Henry³ claims that in cases of cancer of the stomach in which there are no characteristic symptoms present, in which cases it may be difficult to distinguish between pernicious anæmia and cancer, a positive diagnosis may be reached by counting the blood-corpuscles. The author has never seen a case of cancer of the stomach in which the number of red blood-corpuscles has been less than 1,500,000 per cubic millimetre, and has never seen a case of fatal pernicious anæmia in which the number of red cells has not been less than 1,000,-

³ Arch. f. Verdaugshk., April, '98.

000. The reduction of the number of red blood-corpuscles in cancer of the stomach does not keep pace with the cachexia.

Limbeck⁴ has observed that great increase in white blood-corpuscles frequently occurs in cases of medullary sarcoma and lymphosarcoma. The resemblance between the blood in these conditions and in leukæmia is very great, so that a differential diagnosis is not easily made. The increase of white blood-corpuscles in sarcoma is really due to the existence and growth of the tumor, and disappears with the removal of the new growth. Upon this circumstance rests the significance of the examination of the blood in sarcoma. Some years since the left leg of a man who had a medullary sarcoma of the tibia was amputated through the knee-joint. It was desirable to know whether or not there were metastases remaining. The high leucocytosis fell after the operation, and was no greater than could be accounted for by the wound. Some weeks later there was a marked increase of white blood-corpuscles, although neither in the cicatrix nor elsewhere could a new growth be found. The patient failed and died, and the autopsy showed multiple sarcomata in the lungs.

According to Geo. D. Head,⁵ the presence of pus in the abdomen is usually attended by a well-marked leucocytosis. This fact holds true, whether the infected point be localized about a diseased appendix or in a Fallopian tube, or whether the general peritoneum is involved. The height to which such a count may arise varies under different conditions. Sometimes the number of white cells counted may not exceed 15,000 leucocytes, while in other cases the leucocytosis may reach 40,000 to 50,000. In general, the milder the infection, the

lower the leucocyte count, while the more severe the infection, the more white cells will be counted in the circulating blood. The leucocytosis produced by the presence of pus in the abdomen, even though so variable, is usually high enough to be of great clinical value in establishing the presence of a pyæmic infection. However, in some long-standing cases of suppuration where the pus is well walled off from the surrounding tissues, the white count may be normal or only slightly increased. Likewise in very severe, usually fatal septic infections of the abdomen—such, for example, as a general suppurative peritonitis—the leucocyte-count may be normal or only slightly increased. This is the most important exception to the rule of leucocytosis in pyæmic infections of the abdomen, and should always be borne in mind in grave cases of abdominal disease where pus is suspected.

In typhoid fever the leucocyte-count is normal in the first week, while in the second, third, and fourth weeks the count is usually decreased and there is an actual leukopenia. This knowledge furnishes a means of differential diagnosis of the greatest value, particularly in diseases of children. Cabot says: "I have seen good clinicians puzzle for twenty-four hours over a diagnosis between appendicitis and typhoid fever, but the indications of the blood-count were always fulfilled." In a case in which the Widal reaction pointed toward typhoid fever, while the clinical symptoms were those of rabies, the white-blood count was 22,000 corpuscles. The autopsy revealed no typhoid lesions and confirmed the conclusion of the white count.

⁴ Medical News, Jan. 16, '97.

⁵ Northwestern Lancet, July 1, '98.

In tubercular peritonitis there is no increase in the number of white cells in the blood. In sarcoma and carcinoma of the abdominal viscera the white-blood count is of no diagnostic value, since in some cases the leucocytosis is high, while in others little or no leucocytosis exists. The red-blood mixer of a Thoma-Zeiss hæmocytometre and a bottle of $\frac{1}{3}$ -per-cent. glacial-acetic-acid solution in water should be constantly carried in the medical hand-satchel.

Thomas Turnbull, Jr.,⁶ states that in diphtheria there is a marked hyperleucocytosis, which begins early after infection, and is more marked in refractory than susceptible cases. It is greatest at the crisis and steadily decreases during convalescence. In unfavorable cases the leucocytosis continues until death; but in somewhat prolonged cases, with much septic absorption, there may be an uninterrupted decrease of leucocytes continuing to the fatal termination. The staining reaction of the leucocytes is an accurate measure of the severity of the diphtheric infection, and variations in this reaction often precede changes in other symptoms. The increase of poorly-staining leucocytes and amœboid leucocytes usually shows that the disease is taking an unfavorable course. If the leucocytes do not stain better within twelve hours after an injection of antitoxin, it is an unfavorable prognostic sign. Antitoxin, within thirty minutes after its injection, causes a decrease in the number of leucocytes, especially the uninuclear, while the proportion of the well-stained polynuclear cells is increased. In favorable cases the hyperleucocytosis, after the injection of antitoxin, never reaches its original height.

Bacteriological examination of the blood made by Hans Kohn⁷ in pneumonia proved that, in by far the greatest

number of cases which recovered, the cultivation-experiments were negative, whereas in the large majority of fatal cases of pneumococcus was found in the blood. The colonies varied from 2 to 200. The positive results were obtained twenty-four to forty-eight hours before death. The presence of the pneumococcus in the blood is, therefore, of unfavorable import, and these results confirm those obtained by others observers. Thus in a certain number of cases of pneumonia the severity of the disease is due to a complicating sepsis, namely: a pneumococcus sepsis. To the other possible causes of death sepsis must be added. It is possible that the cardiac failure is due to the action of toxins, which may even act more powerfully when the micro-organisms are present in the blood.

G. Lovell Gulland⁸ has discovered a rapid method of fixing and staining blood-films. A small drop of blood, drawn in the usual way, is taken up on the centre of a cover-glass held with forceps, and distributed evenly between that and another cover. The utmost care must be taken to avoid all pressure, as the after-appearance of the red corpuscles depends almost entirely upon the way in which this manœuvre is carried out. The covers are then gently and rapidly slid off one another, and dropped, with the wet side downward, into the fixing solution. This is made up of:—

Absolute alcohol saturated with eosin, 25 c. centimetres.

Pure ether, 25 c. centimetres.

Sublimate in absolute alcohol (2 grammes to 10 c. centimetres), 5 drops (more or less).

⁶ Med. and Surg. Reporter, Jan. 2, '97.

⁷ Deutsche med. Woch., Feb. 25, '97.

⁸ British Medical Journal, March 13, '97.

The quantity required for use at one time, which may be 5 to 10 cubic centimetres for four cover-glasses, should be poured into a wide-mouthed bottle or flat dish, and may be used several times over if it be preserved from evaporation. The writer generally keeps the three liquids in different bottles and makes up the required amount in the above proportions just before using it. The fixation of the elements is practically instantaneous, but the cover-glasses should be allowed to remain in the solution for at least three or four minutes, to fix the film to the cover. They are then taken from the solution by forceps (steel forceps will do), and washed rapidly, but thoroughly, by waving them to and fro in a small basin of water. They are then stained for one minute (not longer) in a saturated watery solution of methylene-blue, and again rapidly washed in water. Next they are quickly dehydrated in absolute alcohol, which at the same time removes the excess of methylene-blue, cleared in xylol, and mounted in xylol-balsam on a slide. The whole process need not occupy more than six or seven minutes; but, on the other hand, any portion of it may be prolonged without injury to the specimen. The fixation may be continued for twenty-four hours, the washing for the same time; but if the staining with methylene-blue be prolonged for more than a minute or two it becomes necessary to use an inconveniently large amount of absolute alcohol to remove the excess of the stain, and the eosin is apt to be washed out at the same time. The red corpuscles are stained pink, nuclei a deep blue, the bodies of the leucocytes in varying shades of pink; the eosinophile and basophile granules in the leucocytes are well brought out; the blood-plates are stained a fainter blue than the

nuclei, and organisms are also well stained. Any other acid stain which is soluble in alcohol and not precipitated by sublimate may be used instead of eosin, and the stain may be omitted from the fixative altogether, so that the cover-glass after fixation in the alcohol-ether-sublimate may be stained in any way that is desired. The cover-glasses used must be scrupulously clean; the simplest way of insuring this is to put them for a few minutes in glacial acetic acid, then wash them with plenty of water so as to remove the acid thoroughly, and dry them with a fine handkerchief. A large quantity should be treated at one time and stored up. The thinner the film, the better the fixation. As it is not possible in specimens mounted in balsam by whatever method exactly to reproduce the appearance of red corpuscles as seen in freshly-drawn blood, it is desirable to control balsam preparations by examining fresh blood, or still better by pricking the finger through a drop of 2-per-cent. osmic-acid solution, and examining the preparation thus made as a fluid mount. This is specially desirable where the red corpuscles are altered in shape to a slight extent; greater degrees of poikilocytosis can easily be made out in balsam preparations.

HOT-AIR TREATMENT.

C. H. Frazier⁹ calls attention to an apparatus which has been employed in a series of cases in the University Hospital, where some three hundred baths were given to test its efficiency. It was found to be most satisfactory. The required temperature can be obtained quickly, in from ten to fifteen minutes, and the apparatus is substantially, but simply, con-

⁹ *Annals of Surgery*, Oct., '97.

structed, and involves nothing that can get out of order or require repair. The cases that were treated included acute and chronic articular rheumatism, gonorrhoeal rheumatism, gout, traumatic arthritis, synovitis, tenosynovitis, and fibrous ankylosis.

The method of administering the bath is as follows: The patient's pulse and temperature are first taken and recorded. The limb, first being completely enveloped in a piece of lint, which is wrapped loosely about the part, is then placed in the cylinder. The time allowed for each bath is from three-quarters of an hour to an hour. At intervals of twenty minutes the door of the cylinder is thrown open momentarily to allow of the ingress of a fresh supply of air. If the patient perspires freely, this opportunity is taken advantage of to wipe the limb thoroughly dry. If this precaution is not taken and the limb is allowed to remain bathed with sweat, there is the possibility, if the temperature is exceedingly high, of a superficial burn resulting. This happened in several cases where the precaution was not taken. The degree of temperature employed varies, some patients bearing with perfect comfort a degree of heat which would be extremely painful to others. The average is about 300° F. The frequency with which the baths are given varies with the severity of the case; usually, however, they are administered on every other day.

Certain physiological phenomena follows the application of heat, such as increased arterial tension, elevation of the blood-pressure, dilatation of the lumen of the blood-vessels, diminution of the erythrocytes, decrease of hæmoglobin, increase in the elimination of nitrogen, and increase in frequency of the heart's action. In cases in which there is a diathesis, either rheumatic or tuberculous,

this treatment can have no beneficial constitutional effect.

Permanent cures of local lesions, symptomatic of diathetic diseases, are not to be looked for from the employment of hot-air baths, but for the relief of joint affections of traumatic origin this method of treatment is most useful and sometimes indispensable, and the results obtained can be called permanent.

E. L. Morse¹⁰ reports a case of malignant senile gangrene cured by topical application of intense dry hot air. The patient, aged 79 years, was attacked with gangrene (senile) in the left foot. Physical examination of the two middle toes, which were affected, revealed a deep ulcerative process, involving also the adjacent structures of the dorsal side of the foot; aside from the general sloughing, the toes were practically dead. Poultices, cauterizations, and antiseptics proved of no avail. Amputation was out of the question on account of the existing heart-trouble (mitral insufficiency), which was quite pronounced. Then decided symptoms of blood-poisoning set in. Added to this the general weakness, due to disease and age, and it was evident that any harsh surgical procedure would almost to a certainty be followed by shock and death. The sloughed tissue was dissected away and the wound treated antiseptically. The condition assumed a worse form, the leg becoming endematous from the knee down and hyperæsthetic. As a last resort, the writer determined to try hot air. The limb up to the lower third of the thigh was wrapped in a towel and put into the cylinder and the air within heated gradually to 350° F. Two treatments a day were given, the temperature reaching later 400° F., for two weeks. Continu-

¹⁰ Indian Lancet, July 1, '98.

ous observation proved a decided diminution of the œdema, healing by granulation of the sloughed tissues, and improvement of the constitutional septicæmic symptoms. Patient was discharged in two weeks, and is working on his farm.

A. Graham Reed¹¹ says a year's experience with the Sprague Hot-Air Therapeutic Apparatus has demonstrated that its use is by no means confined to the treatment of what is generally understood by the terms "rheumatism" and "gout." Its successful application covers a wide range of diseases, particularly of bronchial and asthmatic difficulties, general neurotic conditions, tonsillitis, conjunctivitis, etc., while it is almost a sovereign remedy for sprains or bruises and synovial effusions. Nor has it often been disappointing in its action in the usual types of gout or rheumatism. Even where topi have formed, the solidifications are frequently softened and carried off through the excretory organs.

The skin and kidneys are stimulated by the hot blood, circulation is restored to the affected part, sleep returns to the sleepless, and the general economy is rejuvenated.

In chest and pelvic diseases the patient is put into the body-machine, an apparatus consisting of a metal or treatment section about 30 inches long, with a closed canvas extension for the feet and a canvas curtain at the free end, from which the head protrudes; it being comfortably pillowed, a pleasant sleepiness is frequently experienced by the patient. The heat is diffused over a larger surface of metal than in the leg-machine, and is showered down through a great number of minute openings on to the patient's body, which is covered by a bath-robe. In this machine a temperature of 250° or 260° F. seems sufficiently high to

obtain desired results. Perspiration is profuse, the lymphatic circulation is stimulated, and the joints and ligaments become more flexible.

Age does not seem to be much of a hindrance to recovery; for instance, a Mrs. S., over 80 years old, had rheumatism for the last twenty-five or thirty years, but was very much worse during the last twelve months. Extensive topi had developed in the knees, and she had not been down to her meals for five years. After four treatments the "creakiness" disappeared, and on the day of her sixteenth hot-air bath she took a walk of seven squares. She was discharged perfectly well and apparently a much younger woman. An average temperature of 320° was employed in her case.

As regards the effect on chest affections, a remarkable case was that of a lady with lobular pneumonia of the right lung. After several days of distressing cough, she was placed, on March 20th, in the body-machine for an hour. The congested and inflamed lung was immediately relieved, the cough suddenly left, and up to June 1st had not reappeared. Chronic bronchitis has also been wonderfully responsive to the treatment.

The blood becomes heated from 1° to 5° F., and this seems to be the therapeutic factor. The heat stimulates to action the clogged-up vessels and congested tissues and incites healthy metabolism. Profuse diaphoresis is promoted without the unpleasant head-symptoms usual in a steam-bath, as the patient breathes the ordinary air of the room. The dilatation of the blood-vessels and the flow of blood in the skin are greatly increased, and we have every reason to believe that the same hyperæmia exists in the subcutaneous tissues. After re-

¹¹ Philadelphia Polyclinic, August 6, '98.

moval from the machine in which the patient is usually kept one hour, another hour or more if necessary is allowed for the drying off, the contraction of the skin, the falling of the temperature to the normal, and to the reclothing for exposure to the out-door air.

All cases, as far as heard from, have kept what they gained, excepting in so far as they have returned to errors of diet and lack of exercise. As a matter of course, the originating causes may induce a return of the trouble. Very frequently when a backslider feels an admonitory symptom, he returns for a prophylactic bath.

Sciatica of several months' standing, lumbago, and torticollis have been cured in one application, but there are forms of rheumatism and goutiness which require many and frequently-repeated treatments, taxing the utmost skill of the physician and the perseverance of the patient. Yet, by repeated and well-directed efforts, the soreness is nearly always relieved, and the pliability of the affected parts restored in some degree.

Several cases of rheumatoid arthritis have been treated with more or less success; more, as regards alleviation of the soreness and general symptoms, but less as regards restoration of mobility to the joints involved. In all instances where persistent effort is made, progress of the disease has been checked and relief from acute suffering obtained.

A professor in a Western college was a victim of arthritis deformans, and came for treatment last June. He was obliged to sit with his legs straightened out, and for a year had been unable to feed or dress himself, and could not rise from a chair without help. He was treated in the leg-machine at a temperature varying from 280° to 340°. The diaphoresis

was excessive; his temperature would rise from 2° to 4° during the hour, and hot blood coursed through the swelled limbs, reducing the swelling and relieved the pain. After twelve baths he was able to feed himself and ride a bicycle six miles. After twenty-five treatments he was so much improved that he went north to the Adirondacks, where he carried out various exercises prescribed for him, such as chopping and sawing wood, pitching quoits, etc. The improvement continues to the present time.

In the course of the 1400 treatments given during the last twelve months many exceedingly interesting types of disease have been examined and the great majority have been relieved. The failure of an apparatus to run to a very high temperature must certainly curtail its usefulness. Not that this mode of treatment is a panacea, by any means, but that it is proved to be far-reaching in its direct and correlative tendencies. Acting as well on deep-seated tissues and internal organs as on the periphery of the body, it becomes a most useful adjunct to medical and surgical treatment.

S. Solis-Cohen¹² says that there is no question that it is to be considered among our useful therapeutic measures, and the more one works with it, the more enthusiastic he is likely to be from the results accomplished and the greater is the authority with which he can speak of the range of its application. So far as rheumatoid arthritis is concerned, the author has become satisfied that in his own hands the treatment is almost useless. It is fair to say that the hot air was applied locally only; he did not use the Turkish bath, or Dr. Reed's modification of the Turkish bath, by means of what he calls the "body-instrument." It

¹² Philadelphia Polyclinic, July 30, '98.

is true that pain was relieved, and for a while autohypnotic suggestion made the patients believe that they were improving generally; so that if the patient had been discharged after ten or twelve applications, and the patient herself allowed to report as to the result of treatment, that report would have been extremely favorable. But he continued long enough to enable him to form a judgment as to whether pathological conditions were altered, and he became satisfied that they were not.

J. T. Rugh¹³ has generally observed very good results following applications of hot air for about an hour, the temperature being brought up to about 250°, and in some cases to 300°. The structures about the joints are much softened by such treatment, and yield to forced stretching. The writer has recently corrected the deformity in the majority of a series of cases of flat-foot by baking them thoroughly and applying, under an anæsthetic, a screw form of corrector, and thereby forcibly reducing the existing contractions.

Alice M. Seabrooke,¹⁴ who has had a liberal experience in the use of this agent, says that it is almost always followed by good results, there being only one case in which there was much inflammation about the joints and in which success did not attend the treatment.

STOMACH, DILATATION OF.

Symptoms.—William H. Broadbent¹⁵ states that the imperfect assimilation of food when the stomach is much dilated leads to loss of flesh and strength, and the skin is usually harsh and the complexion pale and sallow. Other symptoms are constipation, thirst, and a flabby or coated tongue. Others, again, are attributable to the effects of ptomaine poisons, generated by septic organisms

in the stomach or intestinal canal, upon the nervous system—headache, depression, which may amount to melancholia, irritability, prostration of strength, and incapacity for mental or physical exertion, occasionally stupor, or even coma.

Nettle-rash and other cutaneous eruptions may be distinctly due to dilatation of the stomach produced by reflex irritation transmitted to cutaneous nerves, or by absorbed matters which affect the nutrition of the skin.

The most characteristic, but by no means the most common, symptom of dilatation of the stomach is copious vomiting daily or at irregular intervals. The time at which it most commonly occurs is in the night, either soon after lying down or in the early hours of the morning, but it may take place at any time. The amount brought up may be astonishingly large, and may seem to be more than the patient can have taken for the previous twenty-four hours. The vomited matters are mostly brown, acid, and evidently in a state of fermentation, but sometimes food taken two or three meals previously may be recognizable. On the other hand, a patient may be sick immediately after taking milk or food, and the vomited matters exhibit no trace of what has just been swallowed, this having been lost in the large amount of fluid which the stomach already contained. The vomiting may be preceded by pain, sometimes severe, and by oppression, but very frequently the nausea and deadly faintness which commonly attend sickness will be entirely absent; the contents of the stomach are easily and rapidly ejected, the discomfort and sense of distension is relieved, and the patient feels well.

¹³ Philadelphia Polyclinic, July 30, '98.

¹⁴ *Ibid.*

¹⁵ Practitioner, Jan., '98.

Very commonly, in cases characterized by vomiting, visible peristalsis of the stomach is started by manipulation, and can be seen to travel from left to right. This affords the best possible demonstration of the size and situation of the stomach.

Palpitation of the heart is very frequent, and is one of the symptoms which compels patients to seek medical advice.

Irregular action of the heart, occasional or persistent, is frequently caused by dilated stomach, as is also intermittent pulse. Like palpitation, the irregular or intermittent action of the heart may occur after a particular meal, or after all meals, and is then most characteristic.

Another symptom is anginoid pain. This, when of gastric origin, is always due to dilatation, or at any rate to distension, and is apparently produced by actual pressure upon and embarrassment of the heart. It may be so severe that, taking only the patient's account of the attack, it could not be distinguished from true angina pectoris, and the extension of pain down the left arm, or down both arms, may be present. The first hint in such a case that the pain is pseudo-anginoid, and not angina of cardiac origin, is often obtained by learning that the early attacks were not necessarily associated with exercise, but that perhaps the very first was extremely acute, and that it occurred in the night or during repose. Angina, unless precipitated by some sudden violent exertion, usually comes on gradually, the pain at first being experienced only when the patient is walking uphill, or against a wind, or goes out on a cold morning, and ceasing with the suspension of the exertion, succeeding attacks being more easily induced and more severe. Gastric pseudo-angina may stimulate true angina

in supervening during exercise, but there is generally more oppression of the breathing than pain.

Another symptom produced by dilatation, or perhaps more frequently by distension of the stomach, is vertigo. The subject of gastric vertigo rarely falls; but the giddiness may be so severe that the patient clings to railings in the street, or makes his way across a room by taking hold of chairs and tables, and it is often excited by movement, sometimes by turning in bed, very frequently by the act of rising from bed. This giddiness appears to be due to mechanical pressure, since it almost always disappears on the escape of a few cubic inches of gas by eructation.

Sleeplessness is a common effect, but a striking characteristic of various symptoms attending habitual distension of the stomach is that they come on after the first sleep, often with remarkable punctuality at a given hour.

S. B. Fowler¹⁶ observed the case of a boy, 16 years old, who, while hauling logs, was taken with pain in the abdomen, and was constipated. Two days later the patient was very weak. Pain in the abdomen, tumor to the right of the umbilicus the size of a fist, and vomiting of faecal matter were present. The abdomen was distended; the lightest touch caused vomiting; splashing in the stomach was heard. Constipation not being relievable, the diagnosis of intussusception was made, and it was decided to open the abdomen. Ether was given, and the patient vomited over a half-gallon of green fluid. Upon opening the abdomen below the umbilicus the first thing seen was the stomach, filling almost the entire abdomen, and almost as thin as tissue-paper. No obstruction

¹⁶ Nashville Jour. Med. and Surg., April, '98.

was found at the pylorus, but, about midway in the ileum, two obstructions were observed in one place. The bowel had invaginated about one and a half inches, and about nine inches from the first place was another, which was invaginated about six inches. The intussusception was overcome, and the patient came out of the operation in good condition. Rectal feeding was given, with no food by the stomach. The boy rested well until the night of the fourth day, when vomiting set in, and he died from collapse. All of the cases reported, so far as the author has been able to see, of acute dilatation of the stomach are seven in number, six of which terminated fatally.

During the course of convalescence from some acute or chronic disease, Fenger¹⁷ claims that the stomach may undergo rapid dilatation.

This condition is marked clinically by a sudden and violent onset; vomiting is violent and intractable: large quantities of fluid are ejected; the fluid is usually greenish, due to admixture of bile. The patient is reduced to a state of relapse or exhaustion, which may prove fatal in a few days.

During the progress of the disease the abdomen becomes distended, the right hypochondrium remaining flatter. The bowels move spontaneously, and a splashing sensation may be elicited over the site of the distension. Sensorium usually cloudy. If treatment is unsuccessful the abdomen becomes more distended, vomiting ceases, and the patient dies of exhaustion.

Indications for treatment are:—

(a) Supportive measures.

(b) Use of stomach-tube one or more times daily, as early in the case as possible.

(c) Rectal feeding.

(d) No food by the mouth until vomiting is nearly or quite controlled. The treatment without lavage is unavailable, and use of narcotics worse than useless.

Diagnosis.—In considering the differential diagnosis Pepper and Stengel¹⁸ say following points must be remembered: Megalogastria may be met with without symptoms of functional disturbance. Such cases can be regarded as physiological or natural largeness of the stomach, provided that there is no sign of stagnation of food, or of disturbance of digestion. Gastropptosis or displacement of the stomach, the vertical position of Kussmaul, or gastropptosis of Glénard, must not be confounded with gastrectasia. In these cases it is most desirable to determine the position of the pylorus and lesser curvature. Obstruction must always be clearly distinguished from atonic gastrectasia. Obstructive dilatation is generally progressive and persistent. The obstructive forms are still further to be differentiated into the malignant and non-malignant.

Etiology.—Pepper and Stengel¹⁹ would classify all cases of dilatation into two groups: (1) *atonic dilatation*, and (2) *obstructive dilatation*. Atonic dilatation may occur (1) by primary or absolute atony of the walls; (2) by secondary or relative weakness. Primary atony occurs in persons of relaxed fibre; in nervous anæmic, or debilitated conditions; or it may be a complication of chronic or even of acute gastritis. Sometimes it may be acute. Relative atony results from overeating and drinking; by adhesions or by dragging of the omentum

¹⁷ Clinical Review, Feb., '98.

¹⁸ American Jour. Med. Sciences, 34-60, Jan., '97.

¹⁹ *Ibid.*

in corpulent persons. Obstructive dilatation is dependent upon pyloric stenosis, and due to diseases of the pylorus itself or to outside causes. Carcinoma and cicatrices of pyloric ulcers are the most important lessons; but, occasionally, other neoplasms or hypertrophy of the pyloric, fibrous, or muscular tissue may be active. Abdominal inspection often reveals abnormal distension. Peristaltic movement can sometimes be detected. The veins of the lower portion of the abdomen, particularly those passing upward over the iliac fossæ, are habitually enlarged and prominent. Succussion-splash can be obtained when the viscus contains fluid. Auscultatory percussion furnishes a reliable indication of the position, and, to a certain extent, of the size of the stomach. In applying the method a double stethoscope with long rubber conductors is used; so that while the patient supports the bell near the position of the fundus, then below and toward the body or pyloric end, the observer is able, without discomfort, to perform the percussion, approaching the stomach gradually from all points. Next, the percussion is reversed, the finger or pleximeter being placed over the stomach itself and near the bell of the stethoscope; then gradually carried outward toward the periphery, until the limits of the organ are passed. Lactic acid, when present, is an indication of stagnation. The amount of urine is another indirect evidence of motor insufficiency.

Treatment.—J. A. Storck²⁰ has used formalin in three cases of dilatation of the stomach with most gratifying results. Where fermentation is marked, the stomach should be washed out before breakfast. Plain water at 40° C. is used first, then 1000 cubic centimetres of a 1-per-1000 formalin solution is now

allowed to flow slowly into the stomach, and withdrawn after two to five minutes, plain water at 40° C. being used finally.

Lyman²¹ says that the first thing to be attempted in gastric dilatation is the emptying of the stomach. This can be most easily and safely accomplished by the use of a soft-rubber stomach-tube, to the upper end of which a funnel is attached. The daily use of the siphon should be continued until satisfactory evidence of improvement is apparent. It may then be introduced every other day, and with less frequency as the case progresses more favorably. The patient should remain in bed for several hours after each operation of lavage, and should apply to the epigastrium and gastric region an ice-bag wrapped in a napkin. A sponge-bath with cold water should be taken every morning on rising and on retiring at night. If so situated that he can use shower-baths of cold water they should be taken every day—preferably in the morning. Light gymnastic exercises and walking in the open air must be enjoined, and the patient must be taught to aim at procuring at least two hours of such exercise—an hour in the morning and another in the evening. If the heart and kidneys are free from disease, horseback-exercise and moderate bicycling will be found useful. The daily application of faradic electricity—placing one pole behind the spleen and moving the other over the region of the stomach for five minutes once or twice a day—is often very beneficial. Good results are also derived from intraventricular faradization,—passing one electrode into the stomach

²⁰ New Orleans Med. and Surg. Jour., March, '97.

²¹ Jour. Amer. Med. Assoc., April 17, '97.

while the other is applied externally in the usual way. The patient must receive his food in small quantities every two hours. Liquids must be largely withdrawn from the diet-lists. Thirst may be relieved by rectal injections of cold water thrown high up into the colon with a long tube. Sugar, starch, and fat should be given in very small quantity. The food should consist chiefly of tender meat well minced, toasted bread, milk, soft-boiled eggs, oysters, and concentrated broths. As improvement appears a larger variety may be gradually introduced. Ten drops of dilute hydrochloric acid should be taken in 4 ounces of hot water after three principal meals each day. One-half hour before each of those meals the patient should take 5 grains of salol or of salicylic acid, or 1 grain of resorcin or of carbolic acid, or 5 grains of sodium hyposulphite, to check the fermentative process during the approaching period of attempted digestion.

Whenever there is pyloric obstruction or such accumulation of contents as to give rise to copious vomiting, Broadbent²² recommends the stomach-tube. The best time for using it is in the evening, when it helps to secure a night's rest.

When lavage is not demanded, and this will be in the great majority of cases, the immediate object will be to relieve existing dilatation or overdistension.

Diet is a matter of the first importance. The meals must not be bulky. Very little liquid, therefore, must be taken with food. Starchy articles of diet must be minimized, both as being bulky for the amount of nourishment they contain and as being liable to fermentation. Only one kind of vegetable should be taken at a meal. Soup need not be absolutely forbidden, but it should be clear

and the quantity should be very small. A small tumbler of hot or cold water may be drunk night and morning, and half an hour before the principal meal.

Overdistension may be actively dealt with by sodii bicarb., 20 grains; sod. sulphocarbonate, $7\frac{1}{2}$ grains; tr. nucis vomic., $7\frac{1}{2}$ to 10 minims, with spt. ammon., tinct. chloroform, and peppermint-water, or other carminative, three times a day, rather nearer the next meal than the last, morning and afternoon and near bed-time. This may be replaced in a few days by an acid tonic with pepsin taken immediately after meals. It is well, however, to continue the evening dose of the alkaline carminative, perhaps without the nuxvomica. Other measures for checking fermentation in the stomach, and obviating or removing distension by gas, are carbolic acid or creasote in 1- or 2-minim doses in pill, capsule, or palatinoïd, given when digestion ought to be completed; creasote with bismuth, charcoal and magnesia, in a cachet, or suspended in a little mucilage. If catarrh has been a prominent feature in the history, bismuth with magnesia or soda, half an hour before food, and hydrochloric acid with strychnine or quinine, or calumba and pepsin, after meals would be indicated. A minute dose of hydrarg. perchlorid., $\frac{1}{32}$ or $\frac{1}{24}$ grain, with the tonic is often of very great service.

When the dilatation is attributable to atony of the muscular coats of the stomach, and is a part of a general neurasthenic condition, arsenic and strychnine are the remedies which have appeared to be of greatest service—say $\frac{1}{24}$ grain of each in a pill with ext. calumb., gentian, or anthemidis, to which may be added

²² Practitioner, Jan., '98.

papain or pepsin, and, if constipation is obstinate, rhubarb or a fractional dose of aloin.

Attention will, in all cases, be paid to the condition of the bowels, and constipation will be obviated by aloetic or other suitable aperients.

Favorable hygienic conditions should be secured as far as practicable.

Boardman Reed ²³ says that the mild-est cases of dilatation of the stomach and some of the worse ones will often yield to diet and exercise alone. The more severe ones, including even those of long standing and those complicated by gastritis, will usually improve greatly upon a careful, non-fermentable diet, lavage, special exercises for strengthening the abdominal muscles and either massage of the abdomen or intragastric electricity, the choice being dependent upon the activity of the secretion of the gastric glands. Massage suits best when there is a deficiency of the gastric juice and is apt to aggravate when there is hyperchlorhydria. This aggravation is especially to be apprehended from a deep, thorough kneading.

Intragastric electricity in the form of a strong faradic current is very efficient in strengthening the weakened gastric muscle and in contracting the dilated organ. Treatment should be given from 5 to 8 minutes at a time on alternate days.

In cases of chronic gastritis with deficient secretion and dilatation, electricity with lavage, massage, and the administration of HCl internally were followed by an increase in the amount of the secretion and contraction of the stomach.

The form of special exercise prescribed for patients with dilatation of the stomach is the use of sets of the now familiar pulleys with elastic cords. No other exercise, unless it be rowing, is so useful

in developing the abdominal and trunk muscles generally.

SYPHILIS.

Symptoms.—D. W. Montgomery²⁴ observes that cerebral symptoms are among the well-recognized phenomena of early secondary syphilis.

Syphilis, according to C. W. Hitchcock,²⁵ figures very largely in the etiology of diseases of the nervous system generally. There are comparatively few forms of insanity in which it does not stand in a causal relation, and one writer has gone so far as to claim that tabes is essentially a syphilitic disease, alleging that fully 90 per cent. of cases of this disease date back to a specific origin. Where syphilis is the prime etiologic factor in nervous affections, its effect almost uniformly is to produce an atypical form of disease. A very long immunity, not only from any nervous manifestations, but also from any secondary or tertiary specific troubles, which may follow the initial lesion, not infrequently renders it difficult for the patient to recall the occurrence of luetic trouble, and, therefore, it is likely to be a common source of confusion and difficulty to the physician. It must always be remembered that syphilis of the nervous system may develop at any time, from a few months to thirty years after the primary infection.

Gumma and arterial disease are, perhaps, the more common forms of the syphilitic lesion. The site of the gummatous process is all-important, and may have much to do with the character of the symptoms developed, whether occlusive, merely mental, or paralytic in character.

²³ Jour. Amer. Med. Assoc., July 30, '98.

²⁴ Medical News, March 6, '97.

²⁵ Medical News, August 6, '98.

The symptoms naturally depend upon the pathological phase present in any given case. While a chronic process may have long preceded the onset of serious symptoms, these may seem to come from a clear sky: *e.g.*, an apparently-acute meningitis may rapidly develop, with violent convulsions, and possibly hemiplegia, more or less complete, follow.

Rubino²⁶ remarks that epileptic seizures may occur in secondary syphilis as well as in tertiary, and states that they are due to lesions of the superficial structures, or to the direct action of the syphilitic virus on the brain-centres, manifesting itself in patients liable to neuropathic symptoms. It is right to regard secondary syphilis as the cause of the epilepsy when this occurs in adult age without any other known cause, and in persons in whom the specific affection has either remained latent or has only given rise to slight symptoms. As regards clinical symptoms, syphilitic epilepsy does not differ from ordinary epilepsy, but differs in yielding to anti-syphilitic treatment. This treatment should be continued for a considerable time.

Legrain²⁷ reports three cases of syphilitic disease of the uterus. 1. The first case, aged 50, had a uterus enlarged to the size of a seven-month foetal head, the fundus being three fingers' breadth above the pubis. The cervix was not discernible. The patient was cachetic, and suffered from greenish-yellow discharge. No ulceration. Antisyphilitic treatment brought the uterus to its normal condition. 2. The second case, aged 54, suffered from metrorrhagia, resulting in anæmia. The uterus was uniformly enlarged, but not fixed. Cervix normal; gumma on tibia. Under iodides the uterus resumed its normal size, the metrorrhagia ceased, and the anæmia

improved. 3. The third case, aged 38, was diagnosed as one of interstitial fibroids. Papular syphilide on knees and elbows. Treated by calomel injections of $\frac{1}{3}$ grain daily. Under a mixed intermittent treatment of protoiodide pills and iodide of potassium the uterus soon became normal.

The author considers these cases to be parenchymatous and fungous metritis, due to diffuse cell-infiltration of the uterus rather than gummata.

Eugene Fuller²⁸ says that primary gumma of the epididymis is exceedingly rare, but late syphilis in the epididymis, not in the form of gumma, is occasionally seen. It is an infiltrating lesion of the epididymis, which generally extends into the parietal layer, giving rise to the feel of a "clam-shell" configuration. This condition is almost always associated with hydrocele.

J. A. Fordyce²⁹ claims that gummatous infiltrations and interstitial inflammations of the kidneys are the result of old-standing syphilis. It is, however, still a moot point whether a specific parenchymatous nephritis is ever the direct result of syphilis in its early stages, for mercury is known to occasion both albuminuria and cylindruria. Aside from the presence of gummata and localized interstitial changes in the kidneys, there is as little characteristic in lesions of these organs produced by the disease as in the symptoms presented during lifetime.

W. Morres³⁰ reports the case of a 50-year-old woman without a specific his-

²⁶ Riv. d'Igiene e Med. Prat., April, '98.

²⁷ Annales des Mal. des Organes Genito-Urin., April, '98.

²⁸ N. Y. Med. Jour., August 6, '98.

²⁹ Jour. Cut. and Genito-Urin. Dis., p. 151, April, '97.

³⁰ Indiana Med. Jour., July, '97.

tory who consulted him for vesical hæmorrhage of a month's duration and growing more profuse. She had lost thirty pounds in weight and was unable to attend to household duties. Physical examination elicited nothing specially satisfactory except as to "saddle-nose" and old bronchitis. The urethra was patulous and in the left upper quadrant of the bladder was a point of special tenderness. The cystoscope showed the urine flowing into the viscus clear. Urinalysis showed no casts, the lesion evidently occupying the bladder. Various measures were tried to arrest the hæmorrhage, but relief was only temporary. After about a month a mass of tissue of the color and consistency of a small oyster was passed from the bladder with great pain. This, on examination, showed small round cells and fibrous tissue, but no evidence of papillomatous growth. The writer was satisfied he had to deal with a gumma of the bladder, and ordered Hardaway's mixture. The hæmorrhage rapidly subsided, but with any cessation of the medicine it would return, with pain at the close of micturition, as before. All urinary discomfort ceased after about two months.

M. Dieulafoy³¹ says that syphilis of the stomach is not so rare as might be believed. The lesions present themselves under various forms; hæmorrhagic erosions, ecchymosis of the mucous membrane, gummous patches, etc. When the symptoms of ulcer simplex are observed in a syphilitic person it is natural to suppose the gastric lesion is of itself syphilitic. It should never be forgotten when in presence of gastric ulcer to look for syphilis in the antecedents of the patient and apply the treatment accordingly.

In syphilitic disease of the heart-wall Sidney Phillips³² says that (1) syphilis may produce gummata or general fibroid

change. 2. Gummata in the left ventricle, except of very small size, are dangerous to life, and when near the apex of the left ventricle may cause sudden death. 3. Gummata in the left ventricle may be suspected if in an individual with syphilitic antecedents there occur signs of derangement of the action of the left ventricle with symptoms of defective or embarrassed action, especially angina pectoris, tachycardia, syncopal or epileptiform attacks; or in the right ventricle when dyspnoea otherwise unaccountable occurs. 4. Extreme feebleness of the heart without dilatation, gradually increasing, in young or middle-aged persons who have had syphilis, suggests syphilitic disease on the right side of the heart. 5. Dilatation of either side of the heart in syphilitic persons may result from syphilitic fibrosis. 6. Hypertrophy of the heart without ascertainable cause and without corresponding increase in strength of heart suggests syphilis. 7. Aneurism of the heart-wall may result from syphilitic local lesions, and may be fatal with or without rupturing. 8. It is probable that gummata and fibroid disease in early stages may be relieved or cured by the usual specific treatment. There can be little doubt that syphilitic heart affections must affect the administration of anæsthetics, more especially as a life-long disease like syphilis is apt to produce local conditions that not infrequently require operative interference. Dr. Kirk found that chloroform is liable to give dangerous results in syphilitic subjects.

On hereditary syphilis W. W. Whitaker³³ remarks that the naso-pharyngeal mucous membrane is ever ready to be-

³¹ Canada Lancet, August, '98.

³² Lancet, Jan. 23, '97.

³³ Laryngoscope, Sept., '97.

come hyperæmic and respond quickly to such infections as pertussis, measles, scarlatina, etc. The fauces in primary syphilis present a characteristic redness. In hereditary syphilis, the same condition obtains, although less actively. The mucous membrane in children predisposed to congestion becomes infected quickly. The lymphatics bearing the infectious agents are stimulated, and the lymph-follicles receive more blood and become active in proliferation. Lymphatic hyperplasia becomes more general and the mucous membrane is invaded. The histological muciparous glands hypertrophy, and increased mucus results. The parts being congested, there is increased local temperature, which, together with the abnormal quantity of mucus, favors self-proliferation and degeneration. When these conditions are established early in life one has the "lymphatic diathesis," to which so much is owed for the many subsequent pathological conditions found in the throat. Gummatous deposits occur in the lymphatic bodies about the muciparous glands in the fauces, under the soft parts. Owing to the liberal distribution of these bodies, extensive areas may be invaded before the process of softening begins, which, when once established, spreads rapidly and causes extensive destruction. Throat symptoms may be anticipated, but seldom encountered until after the age of 5 years. Fortunately inherited laryngeal syphilis is not frequently encountered; but when it does come it is attended with grave conditions. In the very young the tendency to cedema, or laryngismus, may be brought into activity by exposure to cold or some other exciting influences.

V. Dühring³⁴ states that healthy children born of syphilitic parents either possess an immunity against syphilis of shorter or longer duration or show para-

syphilitic manifestations, such as defective development, malformation, etc., or develop about puberty the syphilitic lesions known as late genuine hereditary syphilis. This immunity is known not to be permanent. A set of cases were observed in which there existed proof of hereditary syphilis in the third generation. In malignant syphilis tertiary manifestations are often seen in the early stages of the disease; that is, along with a severe infection there is a pronounced intoxication. This explains how tertiary syphilis may appear to be infective. The appearance of hereditary syphilis in the third generation is due to malignancy.

Diagnosis.—C. F. Marshall³⁵ says many apparently "soft sores" are followed by secondary syphilis, especially in females, among whom the typical indurated chancre is much less common than in males. Again, a soft sore appearing a few days after connection may, after a few weeks, become indurated, owing to double inoculation. Distinctly hard and enlarged glands combined with much induration of the primary sore are almost diagnostic, but the so-called "shotty glands" are very deceptive, and may be felt in many persons after herpes preputialis, or simple inflammation of the smegma-glands of the penis.

According to Marshall,³⁶ it sometimes happens that a patient with some simple skin affection, such as eczema or psoriasis, contracts syphilis. In such cases the resulting secondary syphilide becomes mixed with and modifies the nature of the simple skin affection, so much so that diagnosis is often difficult, and depends mainly on the results of treatment. A mixed eruption of this kind will become

³⁴ Deutsche med. Woch., March 25, '97.

³⁵ Treatment, May 27, '97.

³⁶ Treatment, July 28, '98.

differentiated under antisyphilitic treatment, bringing to view the simple skin affection.

E. G. Janeway³⁷ has observed a number of cases showing danger of error in diagnosis between chronic syphilitic fever and tuberculosis. One was that of a young man who had been sent to a sanitarium for consumption. He continued to grow worse and an examination by the author revealed syphilis. No tuberculous symptoms could be found, and hepatitis had been the cause of his ill health. He promptly recovered under appropriate treatment.

Etiology.—Winkler,³⁸ while examining microscopically the sperm of a syphilitic patient, has observed round, peculiar elements which are not to be found in the sperm of healthy subjects. Wondering whether these elements are not connected with syphilis, he has examined a certain number of syphilitic neoformations and of ganglions, by means of a special method of staining (formol, carbolfuchsin); he obtained the same result. Between the leucocytes in his preparations have been seen some round elements, stained deep violet red, the diameter of which is about a third that of a white blood-cell. They sometimes form groups of two; more often they are isolated. They do not appear to be nucleated. In normal tissues the author has found nothing of the kind.

Niessen³⁹ has obtained from syphilitic tissues, and especially from the blood, pure culture micro-organisms which, when inoculated into animals, present a reaction to human syphilis. When these cultures were injected into the veins of pigs, or inserted subcutaneously, there developed at the point of injection a hard inactive sore, and eight or ten days after the injection there appeared on the skin of the animal numerous bright-red spots

which disappeared after about a week. Rabbits developed at the site of injection similar hard sores. Two of these rabbits were paired, and the female gave birth to a litter of seven, all dead, and two of them, being macerated, greatly resembled in this respect syphilitic human embryos. The author obtained his cultures especially from the marrow and epiphyseal lines of the bones of children who had died from hereditary syphilis. The material was preserved in bouillon, and then grown in various media. He found in almost every instance a variety of streptobacilli or streptococci which he had previously obtained from the blood of patients suffering from dementia paralytica and tabes syphilitica. The bacilli can best be obtained from the blood, after the administration of mercury for a short time during the tertiary period. In the secondary period he had less success in obtaining germs from the blood, which he thought was due to the fact that at this period they lie chiefly in the skin.

J. G. Adami⁴⁰ claims that it is not necessary to have any recognizable first stage of cutaneous chancre. In the female the absence of any superficial or recognizable first stage is especially noticeable; time after time the disease only manifests itself in the secondary stage. The "fixed idea" that there must be a chancre developed at the region of primary infection has led to a thorough and general misunderstanding as to the nature of congenital syphilis. It is a popular fallacy to regard a considerable number of cases, in which the father of

³⁷ Maryland Med. Jour., vol. xxxix, No. 6, p. 574.

³⁸ Wiener klin. Woch., No. 17, '97.

³⁹ Centralb. f. innere Med., May 7, '98.

⁴⁰ Canadian Bract., July, '98.

syphilitic offspring is syphilitic and the mother is apparently free from the disease, as due to the sperm being syphilized, or, if this view be carried to its logical conclusion, it is supposed that the spermatozoön bears with it the syphilitic virus, be it bacillus or whatever the nature of the specific microbe, and introduces it into the ovum at the moment of conception, and thus the offspring develops, syphilized from the start, the mother being and remaining absolutely free from taint. But it is incredible that the germ gain entrance into the spermatozoön, for the spermatozoön being nucleus and flagellum, and scarce anything more, has not the means of ingesting foreign bodies, while we have not a shred of evidence that the syphilitic germ is amœboid and capable of making its way into the spermatozoön. If the syphilitic virus gained entry into the unsegmented human ovum, its effects would surely be to lead to the destruction of the ovum. Foetal syphilis must originate at a later date, and although syphilis in the parents may doubtless have its effects upon the ovum and spermatozoa of the same, and lead to constitutional disturbances in the offspring, progressive syphilitic lesions, the true syphilomata, in the foetus and infant are *not* inherited, but are congenital; that is to say, acquired in utero after conception. If the mother be without sign of syphilis, and the child be syphilitic, the only satisfactory explanation is that the syphilitic virus has entered into the maternal organism and tissues, and has failed to induce any characteristic lesion at the point of entry, but has, nevertheless, through the placenta and chorionic villi gained an entrance into the foetal tissues; the process arrested in the mother has been developed in the susceptible tissues of the child,

and we have here an interesting example of the variability in the manifestations of the disease dependent upon the reactive powers of the tissues. Were any further word necessary in support of this contention it would be found in the significant way in which the liver is affected in congenital syphilis. Extensive specific lesions of the liver in the acquired disease are relatively uncommon. They are the most common of all lesions in the congenital affection. Were the ovum infected it would be difficult to explain why the liver should thus be specially singled out. This organ is the first to receive the blood coming by the umbilical vein; then, if the infection originates from the placenta, hepatic implication is the natural sequence. The essential difference between such congenital, or antenatal, and "acquired," or post-natal, syphilis is that in the former the virus passes immediately into the blood, and so becomes disseminated through the organism, while in the latter the dissemination is delayed. The second stage of acquired syphilis is the first stage of the congenital disease.

M. Delore⁴¹ presented to the Medical Society of Lyons a placenta from a seven months' pregnancy, weighing nine hundred grammes, while the child weighed sixteen hundred. It showed syphilitic gummata, and a greater number of the villosities were obliterated. The child showed superficially confluent petechiæ.

Prognosis.—Krauss⁴² believes that specific meningo-encephalitis and gummata, if small and located in the meninges, offer a very fair prognosis, dependent upon the thoroughness and audacity of the treatment; while complete or even partial recovery is extremely doubtful in

⁴¹ Lyon Méd., July 10, '98.

⁴² Alienist and Neurol., Jan., '97.

all cases where the brain-substance has been infringed upon, or where embolism or thrombosis or other sequelæ of endarteritis exist.

Campana⁴³ concludes that it is not easy to define the duration of contagion, but in any case it is probably longer than is commonly taught; even tertiary gummatous manifestations may be contagious at the beginning. The duration is in relation to the duration of vascular irritative manifestations, and in proportion to these. As long as there are any papular exudative manifestations, contagion certainly exists.

Treatment. — Troisfontaines⁴⁴ draws attention to the extreme importance of feeding with sterilized milk in cases of inherited specific disease, more particularly in those cases in which birth takes place before term.

In the treatment of hereditary syphilis in children⁴⁵ the following methods are recommended for a child 6 weeks old: Twenty drops of Van Swieten's liquid (corrosive chloride of mercury, 2 grains; alcohol, 3 drachms; and distilled water, sufficient to make 4 ounces) four times a day, and at the same time inunction with $\frac{1}{2}$ to 2 grammes of mercurial ointment. The inunction must be made in those parts which do not absorb too readily, and which are not too thickly covered with hair. The inunction should be made at night, and the parts covered with cotton. The child should also be bathed twice a week in starch-water. This treatment to be continued without interruption for five or six months, increasing or decreasing the doses only as needed. One-quarter, $\frac{1}{3}$, or $\frac{1}{2}$ teaspoonful of Gilbert's syrup (hydrarg. biniodide, 3 grains; potass. iodid., 102 grains; water, 3 drachms and syrup, q. s. ad. 10 ounces) should be administered four or five times a day in a bottle of water. The

following prescriptions are given for sublimate baths, which should, however, be avoided when ulcerative affections are present:—

℞ Hydrarg. bichlor. corros, 45 grains.
Alcohol, 7 $\frac{1}{2}$ ounces.

Ft. sol.

Sig.: To be aded to a bath; or

℞ Hydrarg. bichlor. corr, 1 drachm.
Ammon. mur., 1 $\frac{1}{2}$ drachms.
Aquæ, 5 pints.

Mix. ft. sol.

Sig.: For a bath of ten minutes' duration (every second day).

When the slightest intestinal irritation is observed the baths are to be omitted; and if diarrhœa continues the mercury by the mouth must also cease. The inunctions should, however, be continued.

Manassein⁴⁶ says that the living uninjured skin of mammals is impermeable for salves with the usual inunction methods. With the usual inunction methods, salves may penetrate to varying depths into the hair follicles.

Buret⁴⁷ draws attention to the value of local application of mercurial ointments in localized chronic cutaneous syphilides.

Cartier⁴⁸ makes use of the following formula: Corrosive mercuric chloride, 1; sodium chloride, 3; distilled water, 20; in dose of nearly 1 grain of the drug, which is injected into the retro-trochanteric region. This has not caused even the smallest abscess, although severe pain may occasionally be met with. Intramuscular injection should be the

⁴³ Il Policlinico, July, '97.

⁴⁴ Jour. de Méd., March 25, '97.

⁴⁵ Pediatrics, August 15, '97.

⁴⁶ Archiv f. Dermat. u. Syphilis, B. 38, H. 3.

⁴⁷ France Med., July 30, '97.

⁴⁸ Revue de Thér. Medico-Chir., No. 11, p. 361, '97.

treatment of choice in that there is introduced into the organism a known amount of the drug in a condition to be fully utilized. The syphilitic manifestations yield rapidly, not only in severe cases, but in facial manifestations. The injections should be repeated every eight days, exceptionally in five, and four injections are sufficient to relieve the patient of contagious lesions, and six or eight for serious conditions.

E. Lang⁴⁹ claims that subcutaneous injections are more reliable than any other method, and can be graduated to the case. The selection of the preparation is of great importance and requires discrimination, but the chief point is to limit the mercury and not administer too much. The mercurialization must be mild. Inunctions and injections with full doses do more harm than good. The eruptions may be postponed, but the later manifestations are inevitably more severe and prove unusually obstinate. A mild preventative mercurial treatment, combined with a rational mode of life and observation of hygienic measures, is one of the most precious therapeutic measures in our possession.

According to Dabney,⁵⁰ the *modus operandi* of treatment of syphilis by the hypodermic injection of bichloride of mercury is as follows: First, care must be taken to avoid all joints, glands, and blood-vessels. The best sites for the injection are the chest-walls, back, gluteal region, and the upper and outer portions of the arms. The site chosen should be well soaped (green soap) and washed and dried, then rubbed briskly for a few seconds with alcohol. The needle and syringe should be immersed in hot water before each injection. The needle should then be plunged quickly through the skin and well into the cellular tissue, care being taken to avoid invading adipose or

muscular tissue (though some authorities advise deep muscular injection). The injection should be very slowly made, care being taken not to use too much force. As soon as the bichloride solution comes in contact with the albuminous serum, the soluble mercuric chloride immediately becomes the albuminate, which is not very soluble, and herein lies the efficacy and safety of these large doses of this highly-corrosive poison. One of the most striking effects of the hypodermic injection of the bichloride is the way it tones up old, broken-down syphilitics. This treatment does not entirely do away with the iodide of potash, which may be used in conjunction with the injections, especially where the throat, brain, or spine are involved. Wherever mucous patches occur the solid argentic nitrate should be applied.

Lukasiewicz⁵¹ gives the results of treatment of five hundred cases by intramuscular injection of a 5-per-cent. solution of perchloride of mercury, 1 centimetre being injected into the gluteal muscles once a week. On an average, six injections were used for each patient. The cases included all stages of syphilis in both males and females, varying in age from 16 to 65 years. Severe secondary lesions, including rupia, iritis, periostitis, choroidoretinitis, and laryngitis yield to this form of treatment, also early tertiary conditions, such as cutaneous gumata. A peculiar phenomenon appears after the first injection: namely, an increased distinctness of the syphilitic rash, and the author compares this to the reaction which follows the injection of Koch's tuberculin.

⁴⁹ Wiener klin. Rundschau, Jan. 3, '97.

⁵⁰ New Orleans Med. and Surg. Jour., April, '97.

⁵¹ Wien. klin. Woch., April 22, '97.

Whitla⁵² regards mercury as a vital antidote to the syphilitic poison, and so long as the virus of syphilis remains in the organism mercury will expend its force upon it without injury to the patient. This gives a working hypothesis as regards usage. The inutility of iodides in the first and early second stages is emphasized. The continuous method of administering mercury favored by the writer. Small doses should be prescribed as early as possible. Routine treatment deprecated. As a guide determining the effect of the mercury, the weight-chart is strongly recommended. The whole secret of success in the treatment of syphilis is to get as much mercury into the system as possible without producing ill effects.

The general rule for treatment, according to C. F. Marshall,⁵³ should be not to treat any primary sore (in the absence of secondary manifestations) with mercury, unless it is a typically-indurated sore with hard and enlarged glands in the groin.

The best two forms of treatment during active secondary syphilis are: intramuscular injections and inunction.

The best form of mercury is the double chloride of mercury ammonia (sal-alembroth). Ten minims of this (containing $\frac{1}{3}$ grain of perchloride of mercury) is injected into the gluteus maximus once a week, by means of a platinum-iridium needle.

The best preparation to use for inunction is the unguentum hydrarg. cinereum (hydrarg., 1 part; olive-oil, $\frac{1}{2}$ part). About $\frac{1}{2}$ drachm of this is rubbed daily into the skin. The rubbing should be continued for twenty minutes at a time, and different parts of the body used alternately. A warm bath should be taken before each inunction.

Whichever treatment is used, it should

be continued till all symptoms have disappeared. After this it is sufficient for the patient to take mercury in the form of pills, either a grain of hydrarg. cum creta ter die, or 2 grains of pil. hydrarg. bis die, combined with $\frac{1}{8}$ to $\frac{1}{4}$ grain of opium. These should be continued for a further period of twelve to eighteen months, gradually reducing the frequency of the dose. Naphthol (2 to 4 drachms to 1 ounce of zinc oxide) is especially useful in ulcerating condylomata, and also in the treatment of primary ulcerating sores, hard or soft. The best treatment for all tertiary and rupial ulcers is a 2-per-cent. ointment of iodine made with lanolin and vaselin.

Jullien⁵⁴ has observed two cases of secondary syphilis treated by subcutaneous injections of ascitic fluid, taken from a syphilitic patient with cirrhosis of the liver. Precautions were taken to sterilize the fluid. The first patient received 93 cubic centimetres in fifteen injections, with no bad result. However, the effect on the disease was *nil*. The second patient received 520 grammes in 57 injections, increasing from 5 cubic centimetres to 40. All symptoms disappeared, and the patient gained 3 kilos in weight. No further symptoms have occurred up to the present time. No mercury at all was taken at any time.

Gilbert and Fournier⁵⁵ remark that the serum of animals naturally immune to syphilis appears to have no effect upon the course of the disease, and the same is true of the serum of tertiary syphilitics. The serum treatment causes improvement of general condition, disappearance of headache, and of pains in

⁵² British Medical Journal, Sept. 18, '97.

⁵³ Treatment, May 27, '97.

⁵⁴ Semaine Méd., March 16, '98.

⁵⁵ Medical Weekly, iii, 281, '95.

the bones and joints, and attenuation or even disappearance of the cutaneous and mucous lesions, but not constantly or invariably.

Justus,⁵⁶ after examining three hundred patients, concludes that syphilis in the acute stage destroys the hæmoglobin, more or less, and this loss is made up as the disease undergoes spontaneous cure. The same diminution in hæmoglobin occurs after the administration of mercury, and varies according to the amount of mercury employed. The hæmoglobin is restored, sooner or later, according to the severity of the symptoms. It may again sink after repetition of the administration, but if treatment is continued it ultimately reaches a higher level than before treatment was begun. When the hæmoglobin ceases to sink after repetition of the drug, the syphilitic manifestations remit. This sinking after the administration of mercury is a specific phenomenon, and is not observed in the blood of healthy persons, nor in other diseases. This characteristic reaction is observed in early secondary and all subsequent stages. It disappears when the signs of syphilis show remission, but reappears during every relapse.

In a young and vigorous subject with a good digestion, Fournier⁵⁷ recommends ingestion by the mouth as being the simplest and most convenient, unless contra-indicated by other indications. In a dyspeptic subject of feeble digestive powers ingestion by the mouth is not good. One of the other methods should be used. If the patient has bad teeth, with chronically-inflamed gums, inunction is contra-indicated, because it is the remedy most liable to cause stomatitis. In such a case the method least liable to damage the gums is ingestion by the mouth. Social convenience may modify the choice of treatment. For instance,

ingestions should be avoided in people who have to be on their legs most of the day. Again, if the disease is to be kept secret, inunction is out of the question.

Roughly speaking, one may say ingestion for cases of normal syphilis, inunction for severe cases, injection for the worst cases.

An active method is necessary in all cases with grave symptoms, or in specially-malignant types of syphilis. In all cases of iritis, tubercular syphilides, ulceration of the throat, leucoplakia of the tongue, etc., a rapid method is required, viz.: inunction or injection. Injections of calomel are indicated in ulcerating tubercular syphilides and gummata, in phagedenic chancre, in gummatous laryngitis, and in malignant types of the disease.

If the patient has benefited in former manifestations of the disease by a particular method, it is good to repeat this unless contra-indicated in other ways. If other drugs are to be given, such as iron and arsenic for anæmia, or cod-liver-oil, etc., for scrofula, inunctions or injections to be used to avoid the digestive organs.

With regard to the prolonged treatment necessary to combat the syphilitic diathesis, the only good treatment is ingestion by the mouth.

A given manifestation of syphilis will be more amenable to one kind of treatment than to another; this the author names empiricism.

For instance, it is common after several years for syphilitic patients to suffer from slight erosion of the tongue, desquamative glossitis, etc. These cases are very rebellious to inunction or in-

⁵⁶ Brit. Jour. Dermat., Feb. and March, '97.

⁵⁷ Semaine Méd., June 30, '97.

gestion, but clear up rapidly and often permanently under calomel injections. Again, empiricism shows us that inunction is better for hyperplastic glottis, for old tubercular syphilides, tertiary visceral conditions, etc. Inunction is also better in pregnant women and infants.

Colombini and Gerulli⁵⁸ find that iodide of potassium given by the stomach before any other drug, and in the early stage of syphilitic infection, causes an increase in the number of red corpuscles and in the quantity of hæmoglobin. If the administration is continued, sometimes a diminution of the red corpuscles, followed by a progressive and continuous increase, occurs. Sometimes the increase is continuous, and not interrupted by a decrease. When the iodide is stopped the red corpuscles and hæmoglobin tend at first to diminish, but afterward start afresh to increase. At the same time there is a notable and constant increase in body-weight. The authors attribute the beneficial results of treatment to a specific action of the iodide upon the syphilitic virus. They believe that iodide of potassium in moderate doses, and over a moderate time, is the best remedy for the severe forms of chloranæmia due to syphilis.

Cooper⁵⁹ states that he has met at various times a certain number of cases of tertiary syphilitic ulceration which ran a very rapid course, the destruction of tissue being produced by phagedenic inflammation. This phagedenic inflammation cannot be deemed as solely due to the syphilitic poison, and it is very probable that other factors contribute to its production. Where the phagedena attacks the penis, the body, or the limbs, and the face remains free, one of the best modes of treatment consists in the continual warm-bath treatment; but

where the nose is affected this method is obviously inapplicable. The writer has been in the habit of prescribing in these cases a course of Zittmann's treatment, it being the best remedy for arresting the rapid destructive process of syphilitic phagedena. Under this course the destruction of tissue is rapidly stayed, the wound becomes healthy, and cicatrization of it promptly follows. The course of treatment extends to a fortnight, during which time the patient is put upon a strict diet and regimen. The decoctions and pills are made from the following formula:—

ZITTMANN'S DECOCTION, No. 1.

℞ Rad. sarsæ. cont., 4 ounces.
Sem. anisi,
Sem. fœniculi, of each, 1 drachm,
1 scruple.
Fol. senna, 1 ounce.
Rad. glycyrrh. contus., 4 drachms.

And in a linen bag:—

Sacchar. alb.,
Alum. sulph., of each, 2 drachms.
Hydrarg. subchlor., 1 drachm, 1
scruple.
Hydrarg. bisulph, rub., 1 scruple.
Aquæ Cong., 6 pints.

Boil gently down to one gallon, strain, and put into four forty-ounce bottles. Label "The Strong Decoction."

ZITTMANN'S DECOCTION, No. 2.

To the dregs from No. 1 decoction add:—

Rad. sarsæ. cont., 2 ounces.
Cort. lemon.,
Sem. cardam.,
Rad. glycyrrh., of each, 1 drachm.
Aquæ Cong., 6 pints.

⁵⁸ British Medical Journal, June 26, '97.

⁵⁹ Treatment, April 8, '97.

Boil gently down to one gallon, strain, and put into four forty-ounce bottles. Label "The Weak Decoction."

R Hydrarg. subchlor., 2 grains.

Ext. coloc. co., .5 grains.

Ext. hyoscyami, 2 grains.

Ft. pil. 2. Label "The Pills."

The patient is kept in a room at 80° F. The diet consists of: Breakfast—Boiled egg and bacon, tea; no sugar or spices. Lunch—Butcher's meat, vegetables; no fruit. Dinner—Soup, fish, poultry.

The evening before beginning the treatment the two pills are taken, and the next four days, at 9 A.M., 10 A.M., 11 A.M., and 12 M., half a pint of the strong decoction drunk very hot.

At 3 P.M., 4 P.M., 5 P.M., and 6 P.M., half a pint of the weak decoction cold. The patient is kept in bed, except for one hour every evening. On the fifth day he is allowed to get up; he may have a hot bath, and dress, and is allowed, if he asks for it, a little brandy or whisky and soda. In the evening 2 pills are administered, the patient starting the decoctions the next day as before. So the treatment goes on until the fifteenth day, when it is discontinued.

Lyman Ware⁶⁰ has not seen a single case of syphilitic neuroretinitis following thorough and prolonged antisymphilitic treatment. It never terminates in spontaneous recovery. Without treatment it is sure to end fatally. Large doses of potassium iodide are of value in arresting the disease in some severe and dangerous cases, but they do not compare with mercury in eradicating the syphilitic poison. The manner in which the mercury should be used is of secondary importance. For years the author relied mainly on inunctions, but recently has resorted to hypodermic injections of the

preparation proposed by Althaus, which consists of metallic mercury, 1 part; lanolin, 4 parts; carbolic oil (2 per cent.), 5 parts. The usual dose, 5 minims, may be gradually increased to 10, giving an injection once a week, in the region of the glutei muscles.

Fruitnight⁶¹ thinks that if a syphilitic woman becomes pregnant she should at once be placed upon specific treatment, and thus probably secure the escape of the fœtus from the threatened infection. Treatment should be pushed vigorously up to the labor. When it is known that either one or both parents are syphilitic, procreation should be forbidden for a period of at least two years, during which active treatment should be instituted. Then, after six months have elapsed without treatment and with no manifestations of the disease, procreation may be permitted. When there is a reasonable suspicion of the disease in an infant treatment should be begun immediately after birth and vigorously continued. The remedy *par excellence* is mercury. In young infants the best method of treatment is that by inunction. The place of inunction should be changed from day to day. A convenient way of using this method is to place a piece of the ointment on the belly-band, and by the movements of the child and the ensuing friction it is easily rubbed into the skin. The officinal blue ointment is used in this way, but it must be diluted according to the age of the infant. If the symptoms be urgent, or if it be desired to bring the system more rapidly under the influence of the mercury, calomel should be given in doses of $\frac{1}{10}$ of a grain four times daily. The treatment with mercury should be con-

⁶⁰ Archives of Ophth., p. 345, '97.

⁶¹ Medical News, July 30, '98.

tinued for at least one year. During the last half of the year the amount per day can be reduced by one-half or by one-third. If the mercurial treatment be persisted in for a longer period than one year it will be advisable to interrupt its administration occasionally. Salivation is very rare indeed. The symptom which would indicate that the child is becoming profoundly affected by the drug is the occurrence of pronounced anæmia. If the symptoms resemble those of the tertiary stage in the adult the iodide of potassium may be given, from 20 grains to 2 drachms *per diem*, according to the age of the child and the severity of the case, and, of course, largely diluted. In conjunction with the specific treatment, the tonic treatment is of the utmost importance. Among the remedies to be employed cod-liver-oil, iron, arsenic, and nux vomica hold a prominent place. The local lesions of hereditary infantile syphilis will demand the same treatment which is given such lesions in the acquired variety of the disease, whether in the adult or infant, the strength of the remedies employed always being adapted to the age of the patient.

Ramón Guitéras⁶² says that almost every case of common chancre or initial lesion is generally of the form of a simple erosion or ulceration. If the former, the patient is directed to wash it two or three times a day, and to dust on a mild antiseptic powder, consisting of boric acid, bismuth subnitrate, and calomel, equal parts, and afterward to put on a thin layer of absorbent cotton to absorb any discharge. This powder is very satisfactory when the lesion is situated on the glands or on either side of a retractile prepuce or on the body of the organ. If this does not have a curative effect in a few days, a wet dressing

of black-wash should be used. If it is the ulcerating form, it should first be cauterized with a saturated solution of silver nitrate, and then dressed in a similar manner with the powder mentioned. If it is the dry, scalding papule, a wet dressing of black-wash is best. A thin film of cotton spread over the lesion and kept moist by pouring this solution upon it is usually sufficient. In subpreputial lesions the method first to be tried is that of a subpreputial astringent or antiseptic injection. This should be given every three or four hours, after first injecting plain water to cleanse the parts. The materials usually used are a weak solution of carbolic acid (1:250) or of mercuric chloride (1:10,000), or black-wash. If, however, under such treatment, the condition seems to grow worse, a dorsal incision should be made through the entire thickness of the prepuce, and, after thorough bichloride irrigation, the lesion should be cauterized, and then treated by local application of a powder or solution and absorbent cotton, as mentioned. This should be followed later on by a circumcision. In chancre of the meatus a solution seems to produce the best results. *Lotio nigra* should be kept in constant relation to the lesion by means of an absorbent-cotton plug saturated with it, a fresh one to be inserted after each act of micturition. In addition to this, an alkaline solution or a urinary antiseptic should be given internally to render the urine less irritating. Extragenital chancres are treated in the same manner as those on the genitals: *i.e.*, cauterized when necessary, and treated with mild antiseptic or astringent washes or powders. The author does not think it wise to give mercury in the first stage, excepting in

⁶² Indian Lancet, June 6, '98.

certain cases where diagnosis is absolutely sure. It is advisable when the initial lesion may do harm to the patient or to others, as in chancre of the lip, finger, nipple, etc.; when there is extensive tissue-ulceration; or when certain disagreeable symptoms, forerunners of the second stage, have appeared, as intense headache or pains in the bones. The favorite preparation in the second stage is the protiodide. It is given in pill form, the strength usually being $\frac{1}{6}$ grain. A good method of increasing the strength is that advocated by Dr. Keyes, which is practically as follows: 1 t. i. d. for the first three days; 1 night and morning, and 2 at noon for the next 3 days; then 2 night and morning and 1 at noon for 3 days; and afterward t. i. d. for another 3 days; then 7 a day, increasing in the same manner; then 8; then 9; and so on until the symptoms of mercurial saturation begin to show themselves, such as colicky pains, diarrhœa, sore mouth, foul breath, etc., etc. When this point has been reached the same strength may be maintained, and the symptoms relieved by taking, in addition, a small amount of Dover's powder or some other opiate three times a day. If symptoms of salivation occur, it is a sign that the patient has taken too much mercury and it should be stopped. It is always well to instruct the patients to be on their guard against salivation, to keep their teeth clean, and every morning on arising to click them together, and in case the gums or teeth feel sore to immediately leave off the medicine. It is well to have the teeth put in order thoroughly before beginning treatment; to brush them after each meal and on arising and retiring, each brushing to be followed by rinsing out the mouth with listerin or borin diluted with 4 parts of water. The

author does not believe in giving an opiate to counteract the intestinal symptoms of large doses of mercury, as it is likely to create in the patient the opium-habit, and it should never be used until other preparations have been tried which may, perhaps, be tolerated. The patient's condition may often be improved wonderfully in cases of diarrhœa by giving iron in connection with the mercury. Atropine, belladonna, and hyoseyamus also seem to prevent certain symptoms of mercurial poisoning. If the patient cannot stand 9 grains of the protiodide a day, it is much better to try some other preparations. It is well to have the patient on the largest dose that can be agreeably borne, so long as the active symptoms of syphilis are present, and then to fall back to $\frac{3}{4}$ of this dose, and to continue on the $\frac{3}{4}$ allowance until active symptoms reappear, when the patient should again be put upon the full dose and kept upon it during their activity, when the dose can again be dropped to $\frac{3}{4}$. If the protiodide cannot be well borne, the author knows no better salt to try next in order than the tannate. This may be given in pills of $\frac{1}{2}$ to 1 grain in strength, and may at times be increased until the patient is taking 5 grains a day. This form seems to be well tolerated, and the patient obtains more mercury than by taking the protiodide. The pills are ordered as follows:—

R Pil. hydrarg. tannici oxydulat.,
of each, $\frac{1}{2}$ grain.

Sig.: One three times a day, and increased as directed.

Inunctions of mercurial ointment are good for severe cases, and when salts of mercury cannot be well borne when given internally. From 20 to 60 grains may be used at each treatment, and

should be rubbed in as follows: The first day, on the inner side of the legs; the second day, on the inner side of the thighs; the third day, in the iliac regions; the fourth day, over the sides of the chest; the fifth day, over the inside of the upper arms; the sixth day, over the flexor surfaces of the forearms; the seventh day, rest. These should be taken at night before retiring; a bland ointment may be applied after each inunction. Woolen clothing should be worn during a course of inunctions. Symptoms of salivation should be watched for during this treatment, as it is liable to come on very suddenly. Fumigations are very efficacious if one wishes to produce a very rapid effect, especially in late secondary lesions. Here calomel is used, from 20 to 60 grains at each sitting. The patient is seated in a chair, with a covering or blanket extending from his neck over himself and the chair to the floor. The calomel in the pan beneath is then heated, and he is subjected to the fumes for from twenty to thirty minutes. Hot baths have no specific effect on syphilis, but they are good as a measure of hydrotherapy, and mercury by inunction is better absorbed. Occasionally on the forehead there are a number of lenticular papular lesions forming a corona veneris. In this case it will be well to apply the ammoniate-of-mercury ointment on a piece of sheet lint on retiring and to allow it to remain on over night. Ecthymatous, impetiginous, and pustulo-crustaceous syphilides are also benefited by the same applications. Palmar syphilides may also be treated by the white-precipitate ointment alone, or mixed with equal parts of the zinc-oxide or boric-acid ointment. In onychia and paronychia the ammoniate of mercury should be applied locally, and a glove-

finger worn over the parts for protection. Moist papules about the genitals are best treated by the powder of bismuth boric acid and calomel, with a dressing of absorbent cotton to keep them dry. In case of mucous patches in a man's mouth, his tobacco should be cut off, and he should use a mouth-wash of 1:2000 bichloride solution in peppermint-water four or five times a day, in addition to which an application to the patches should be made every three or four days with a one-in-eight solution of silver nitrate, or 4 per cent. chromic-acid solution. The best way of doing this is by twisting a thin film of absorbent cotton about the end of an applicator or a tooth-pick and then moistening it in the solution, when it is applied to the mucous patch until this becomes whitened. In syphilitic alopecia the head should be washed night and morning with a 1:1000 solution of bichloride, or an ointment of ammoniate of mercury and boric acid may be applied. Throughout the second stage the patient should be kept in the best possible health. He should have plenty of plain food, fresh air, and exercise. Stimulants can be indulged in if well tolerated, but should be limited to light wines with the meals, spirit being strictly interdicted. Smoking in moderation may be allowed if it does not irritate the mouth and cause mucous patches. Sexual intercourse should be forbidden until all active symptoms of the disease have disappeared. It is the writer's custom to give mercury for one year and then to change it to mixed treatment, which is continued for another year, $\frac{1}{6}$ grain of biniodide of mercury and from 3 to $7\frac{1}{2}$ of potassium iodide in the compound syrup of sarsaparilla being given. After the two years of treatment have been completed, patients are recommended to

take mixed treatment for a period of six weeks every spring and fall for two or three years. In the secondary stage of syphilis potassium iodide should not be given, excepting in precocious cases in cachectic subjects when dangerous symptoms, such as those of the nervous system depending upon central gummatous deposits, have appeared, or when from similar lesions a destructive process has started in the respiratory or genito-urinary tract or elsewhere. In the ordinary cases of tertiary syphilis good results are obtained by giving the "mixed treatment" internally and using mercurial ointment externally, and it is only

when some serious or active process takes place that it is necessary to give the very large doses of iodide. Potassium iodide, a grain to the drop, may be given in water or milk with the compound sarsaparilla or syrup of orange-peel, and may in some cases be run up to 200 grains a day, which is usually sufficient to control any active tertiary lesion. As high as 800 grains of this salt have been given. Some say that this should be taken on an empty stomach, but the presence of a little food does no harm. The conditions which indicate the use of iodide are an excess of cell-growth and an accumulation of the same.

Cyclopædia of Current Literature.

DIARRHŒA, CHRONIC.

Etiology.—Among the causes of diarrhœa the different conditions of the stomach that give rise to abnormal intestinal fermentation and irritation are of the utmost importance.

The causes of diarrhœa in cases presenting an absence of gastric secretion may be various. One of these causes is the precipitate manner in which the stomach propels its contents into the intestines. Sometimes the stomach is empty one hour after a meal.

Again, the chronic inflammation of the gastric mucosa may extend into the intestines and give rise to diarrhœa from chronic catarrhal enteritis, or the intestine may simply rebel from over-taxation, or the diarrhœa may be the result of intestinal irritation from unusual toxic substances developed in the bowel. The diarrhœa differs in different cases. There are invariably several morning evacuations, and after the middle of the forenoon no further trouble in

some cases; in others, the diarrhœa is post-prandial; in a third class there are many evacuations during the day, taking place irregularly. In some cases there is diarrhœa every day; while in others it is periodic, coming on suddenly, lasting a week or two, and then followed by constipation or by regular daily evacuations. In some cases there is a good deal of intestinal flatulency. Pain is not a common symptom in these diarrhœas. There is usually no elevation of temperature; the temperature is sometimes below normal. The appetite in some cases is voracious. The character of the stool varies. In some instances there is an abnormal amount of mucus. The discharges are frequent and watery. In some cases there may be constipation; but this is less common.

Treatment.—In the treatment of this intestinal affection hydrochloric acid is most efficacious; 20 to 30 drops of the dilute acid may be given after meals, and the dose repeated in an hour. Jaworski

says the acid is needed to develop the formation of pepsinogen in the gastric glandules. The acid is quickly neutralized and combined in these stomachs. In some cases its presence cannot be detected ten minutes after administration. Allen A. Jones (Boston Medical and Surgical Journal, August 4, '98).

DYSMENORRHŒA.

Etiology.—Dysmenorrhœa is caused, in a large number of cases, by a malformation of the uterus, to which must be added the thickening of the mucous membrane along with the congestion, which is natural when moderate in amount, at the time of the menstrual flow.

In a case of dysmenorrhœa with ante-flexion, the bend *per se* is not alone the cause of pain. It is only when congestion, with its resultant thickening of the mucous membrane from simple turgescence, or that change which goes by the name of hyperplasia, supervenes, and we are face to face with a potential or actual organic stricture, that pain becomes a prominent symptom.

There is no pain at the beginning of menstruation beyond the backache, and it is not until one or more years have elapsed that the special pain begins, at first in a minor degree, but gradually increasing up to the age of maturity. The exception to this will be found in the case in which the patient happens to be the subject of catarrh when the menses first make their appearance.

The explanation is as follows: From the very beginning—assuming that there is no catarrh, and consequently no congestion—the flexion causes some retardation of the flow with no more pain than the ordinary backache, or a sense of weight in the pelvis; the frequent repetition leads by degrees to stasis of blood

in the organ, and pain of a special character gradually comes on, yet varying from time to time, probably according to variations in the degree of congestion.

Flexion of the cervix proper causes only a minor degree of dysmenorrhœa, and it is only in cases of flexion at the site of the internal os that we meet with the more severe forms. In these cases the symptoms forcibly remind one of the first stage of labor, which is so often accompanied by vomiting.

The most extreme constriction of the external os does not cause dysmenorrhœa at all approaching in severity to that produced by constriction of the internal os.

Treatment.—Only three rational methods are recognized:—

1. Division of the cervix.
2. The stem-pessary.
3. Dilatation.

The operation involves the dominant idea that there is a stricture to be overcome, and that this stricture affects the internal os. No operation upon the external os can avail anything when the condition to be remedied has its site at the internal os.

The stem-pessary is of value. Its range of application, however, is limited; married life is a bar to its employment, and it can only be used in the unmarried or widowed subject. In several instances of patients contemplating marriage the writer advised the postponement of that proceeding, has caused the patient to wear a stem-pessary for from nine to twelve months, and removed the instrument a week or two before the event. In every case the dysmenorrhœa has been relieved, and the patients have subsequently become pregnant. The presence of a foreign body in the cervical canal causes dilatation of that canal

to an extent much beyond the size of the body that occupies it. Were it not so, the dysmenorrhœa would be aggravated rather than relieved by the treatment.

For the form of antelexion associated with retroversion of the organ there is no other treatment that can be of any avail than that by the compound stem.

No one, who has had any appreciable amount of experience, believes that a vaginal pessary can beneficially affect an antelexion of the uterus.

Gradual dilatation is simple, and is attended with so little danger that it may be done in one's consulting-room. The process is repeated as often as possible at intervals of three or four days before the next period comes on, and, if begun within a few days of the last period, temporary relief may be afforded in one intermenstrual interval. In a considerable proportion of cases it will be necessary, or at least advisable, to repeat the dilatation (at one sitting only) after a lapse of several months. By watching the case in this way a permanent cure may be affected. When congestion is well marked there will often be free bleeding; but this, as a rule, helps to diminish the congestion, with the aid of other means to that end. To secure a good result the dilatation should be repeated until blood is no longer drawn.

In the case of patients from a distance, and in the more severe cases when practicable, especially when congestion is a prominent feature, the patient should be confined to bed, the bougies passed every second or third day, and the instrument left in as long as the patient can bear it.

Rapid dilatation for dysmenorrhœa is not favorably regarded. G. Granville Bantock, (*British Gynæcological Journal*, August, '98).

EUCAINE.

Eucaïne considered the best local anæsthetic. Compared with cocaine, solutions of greater strength are required; but eucaïne is far less poisonous, and stronger solutions may be safely employed. From 6- to 10-per-cent. solutions used, and in over fifty cases not a single instance of toxic symptoms noted. Injections of 5 grains of the hydrochlorate in one sitting is usually sufficient for an operation of considerable magnitude, —*e.g.*, herniotomy,—but as much as 9 grains may be used. The benumbing power of eucaïne is quite as certain as of cocaine, without the prostration with cardiac failure, which often follows cocaine in smaller doses. Eucaïne has a very persistent effect, and on the mucous membranes does not act quite so effectively as cocaine on mere contact. On this account, in operations on tissues covered by mucous membrane, a little cocaine solution may be first applied, and then the eucaïne injected. If the operation is to be a severe one, or prolonged, the patient should be given a small dose of morphine a quarter of an hour before, for the effect on the mind. While the skin is perfectly anæsthetized, the tissues beneath it are not, and section of muscle causes a dull pain not hard to bear. Lilienthal (*Annals of Surgery*, May, '98).

GLANDULAR FEVER IN CHILDREN.

Symptoms.—Acute glandular condition in children observed which has hitherto been unrecognized. The illness commences suddenly in children between the ages of 2 and 4, with coated tongue, vomiting, headache, abdominal pain, and rapid rise of temperature. These symptoms are accompanied by tenderness in the anterior triangle of the neck, and by stiffness and pain on movement

of the head. There is usually pain on swallowing, but there is never, during the whole course of the illness, any pharyngitis or tonsillitis. Generally, two or three of the lymphatic glands, under the anterior edge of the sterno-mastoid, enlarge to the size of a walnut, and are acutely painful on pressure. They remain in this condition for two or three days, during which time the temperature continues to rise, sometimes reaching 104° F. Gradually the glandular enlargement subsides, till, by the end of two or three weeks, complete resolution takes place, suppuration never occurring. The temperature falls by lysis, accompanying the subsidence of the glandular swelling; almost invariably the glands attacked are those on the left side of the neck, but occasionally, after they have subsided, the glands on the right side become affected in an exactly-similar manner. In a large proportion of the cases there is a palpable enlargement of the spleen and liver. The enlargement of the gland is a distinctly idiopathic condition. The disease is sometimes mistaken for mumps on superficial examination, but the swelling in the cases described is confined to the glands under the sterno-mastoid, the parotid gland remaining quite unaffected. Pfeiffer (*Jahrb. f. Kinderh. Leipzig*, B. 29, S. 257, '98).

GOLF.

Golf combines exercise, pleasure, and fresh air without the risk of injury to heart, lungs, or nervous system of some other exercises in which there is high blood-pressure and arterial tension. The game is not contra-indicated in heart-lesions, arterial calcification, albuminuria, old age, childhood, and certain hysterical conditions which would be aggravated by such exercise as bicycling,

swimming, horseback-riding, or by mountain-climbing. In all affections marked by slowing of oxidation, or in those consequent upon intoxication by the products of organic disassimilation, the game of golf is to be recommended as the best adjuvant method of bringing about a cure. The obesity and degeneration of middle age when the biceps has diminished and one's energy is wanting may be helped by devotion to golf. The further tendency of the exercise is to eliminate the so-called diathesis and thus do away with gout, lithæmia, headache, and dyspepsia; while its hygienic and therapeutic consequences are admissible in cardiac and pulmonary affections. Although moderation is advisable in such circumstances, there can be no doubt of the benefit derived in some cases of cough, nervous asthma, and in affections of the bladder and prostate. But it is pre-eminently in functional nervous disease that our great Anglo-Saxon game is to be recommended, both as prophylactic and curative. No exercise or recreation is better for the mentally overworked, the hysterical, the melancholic; none helps to preserve concerted action of the eye, brain and muscle, known as the psychological movement; none, perhaps, with the exception of swimming, gives one so good an appetite; and, as to insomnia, such a thing scarcely exists among the devotees of golf. Irving C. Rosse (*Journal American Medical Association*, August, '98).

HYPERTRICHOSIS.

Treatment. — Hypertrichosis may be successfully relieved by means of the Roentgen rays. In order to avoid the production of inflammation a current is used which does not exceed a maximum of 2 ampères, the maximum tension being 11 1/2 volts, and the source of the

light is placed at a distance of 20 to 25 centimetres from the skin which is to be subjected to the influence of the rays, which are not allowed to act for a longer period than ten minutes. (When it is desirable to produce inflammation for some therapeutic purpose $3\frac{1}{2}$ ampères, $12\frac{3}{4}$ volts, and a distance of 10 centimetres are employed.) The apparatus used is supplied by M. Kohl, in Chemnitz. In seven cases of epilation the best therapeutic results were obtained after seventeen to thirty sittings of short duration. In several of the cases it was noticed that from one to two days elapsed before the hair fell out. In several brunettes the hair became, before it fell out, snow-white. Eduard Schiff and Leopold Freund (Wien. med. Woch., Nos. 22 to 24, '98).

LACTOPHENIN AS HYPNOTIC.

Excellent hypnotic effects obtained with this drug in a long series of cases of insanity of various forms. From 1 to 3 grammes of lactophenin in 150 grammes of gum-arabic solution, taken an hour after the evening meal produced a deep and quiet sleep within half an hour; the patient awoke after from four to nine hours; no unpleasant after-effects were noted. The author states that the drug seems to exert no disturbing influence on the heart's action, and is therefore specially suited for the insane who are subject to cardiac or vascular affections, and lung and kidney diseases. In case of reduced effect through habitual use, a temporary change to some other hypnotic is advised. Christiani (Pharm. Centralhalle, August 25, '98).

MORPHINE POISONING.

Treatment.—Cocaine used in a case of morphine poisoning with success. The trial was made on account of the

knowledge that opium and cocaine act very largely on the nervous centres, but in opposite directions. Thus, opium retards respiration, cocaine stimulates; opium slows the pulse, cocaine accelerates it; opium checks the secretions, except sweat, cocaine increases them; opium contracts, while cocaine dilates the pupils; opium produces sleep, cocaine wakefulness; opium retards peristalsis, cocaine increases it. The same opposing effects are noticed also in their action on the kidneys. The cocaine was given hypodermically in $\frac{1}{4}$ -grain doses, given first with apomorphine and strychnia, and then with small doses of potassium permanganate. The injections were repeated two or three times, and recovery followed by the next day. C. W. Williams (Lancet, vol. xviii, p. 192. '98).

PROSTATITIS, CHRONIC CATARRHAL.

Diagnosis.—In the vast majority of cases catarrhal prostatitis occurs as a result of chronic posterior urethritis, the acini or follicles of the prostate becoming involved in a chronic catarrhal inflammation from extension of the same process from the mucous membrane of the deep urethra.

Among the more prominent clinical features of this affection are: 1. Prostatic discharge at the meatus upon waking in the morning. This condition is frequently mistaken for gleet; as a matter of fact, the discharge is distinctly different from that of true gleet, in that it is colorless, resembling glycerin in appearance, and does not stain the linen. Its presence at the meatus is probably due to an atonic condition of the compressor urethræ muscle, permitting some of the prostatic secretion contained in the follicles of the gland to enter the

anterior urethra. An examination of the first morning urine will positively settle the character of this discharge, the absence of clap-shreds in this urine proving conclusively that the discharge is entirely prostatic in character. 2. Prostatorrhœa: this condition is brought about by the overdistension of the glandular tubules with prostatic secretion. Its occurrence is noticed, as a general thing, at defecation, or at the close of urination, and in some few aggravated cases it appears after some unusual physical exertion. 3. Constant urethral pain, located either at the glans penis or in the perinenum, relieved temporarily by urination.

Another group of symptoms, causing much mental perturbation to these patients, is increased frequency of urination, associated with a forked or sprinkling stream, with perhaps some little dribbling. A rectal examination is essential in all cases where one is led to suspect the presence of catarrhal prostatitis. If the prostate be the seat of a chronic catarrhal inflammation, it will be found to be somewhat larger than normal, and soft to the touch; stripping the gland causes much pain, and is invariably followed by the appearance at the meatus of an abundant prostatic discharge, which upon microscopical examination is found to be made up of granular phosphates containing pus. H. M. Christian (*Jour. Cutan. and Genito-Urin. Dis.*, August, '98).

PROTARGOL.

One of the latest preparations suggested for use in inflammatory affections of the urinary tract, and found by many to be of great service, is a silver combination with proteid—protargol. It is in the form of a fine, light-yellow powder, readily soluble, with a brown color,

in distilled water (1 in 2), the solution being slightly alkaline in reaction, and giving a precipitate with hydrochloric acid, soluble in excess. Protargol is not simply a mixture of proteid and a silver salt, but a true chemical combination. It acts as a strong antiseptic, in 1.5-per-cent. solution, exerting a marked effect upon gonococci, while it does not irritate the tissues. Its mode of action is doubtful, but probably arises from dissociation of the silver salt from the proteid through the action of the bacteria, either from the acids formed by them or by their attacking the proteid moiety to succumb to its silver partner. Editorial (*Edinburgh Medical Journal*, July, '98).

PULMONARY TUBERCULOSIS.

Treatment.—The best beech-wood creasote can be given with benefit, in amounts varying from 120 to 240 minims daily, in cases of pulmonary tuberculosis; the drug is best administered in codliver-oil, or in spirituous solution, and in some cases the "creasote-chamber" or oro-nasal inhaler may be ordered in addition, with advantage. The dose should be small at first, but it can be rapidly increased to 40 minims, three times daily, for an adult. Large doses rarely cause any gastric disturbance; on the contrary, the appetite is frequently increased. The cough, expectoration, and night-sweats are diminished, and the physical signs improved. Owing to its disinfectant action in the alimentary canal, the drug probably diminishes the risk of tubercular enteritis by autoinfection when patients swallow their sputa; but, owing to the increased peristalsis, which is created by creasote, it is usually contra-indicated in cases where the ulceration is already advanced. The drug does not tend to cause hæmoptysis, but rather to prevent its recurrence.

Creasote does not irritate the normal mucous membrane of the genito-urinary tract. Tamplough (British Medical Journal, May 28, '98).

RECTAL PROLAPSE.

Treatment.—Child suffering from debility and chronic diarrhoea developed rectal prolapse, which, during twenty-four hours, increased to 10 centimetres. On account of complete paralysis of the sphincter and impossibility of retaining the reduced gut in position, the anus was temporarily closed with three catgut sutures and the diarrhoea treated with proper remedies. On the third day the sutures were removed, which action was followed by a slight prolapse. Plaster strips were thereupon substituted for the sutures, and later radial cauterizations of the anal mucosa resorted to. Complete recovery of child followed. O. Helms (Hospitaltidende, 1898, No. 24). [Report of Corr. Ed. Johnson, Copenhagen.]

SEROUS PLEURISY.

Treatment.—Eleven cases of serous pleurisy treated with a mixture of guaiacol, 1 part, and tincture of iodine, 4 parts, with favorable results. A drachm of this mixture was applied once daily to the affected side, which was then covered with wax-paper, cotton, and then with a bandage. Besides this treatment, the patients received only small doses of codeine of Dover's powder. In all cases the exudate became absorbed more quickly than was observed by the author under any other method of treatment. Neither objectively nor subjectively were there any disagreeable by-effects noticed; the irrigation of the skin was also insignificant. From five to seven guaiacol applications were required to cause a complete disappearance of the exudation. The

SORE NIPPLES, THE PREVENTION OF.

temperature was taken before and after each application, and a fall of 0.4° to 3.5° F. was recorded. In cases with normal temperature the reduction was insignificant. The author believes that by irritating the peripheral nerve-endings the guaiacol acts on the thermal and vasomotor centres; hence the reduction in temperature and increased absorption-power of the pleura. Besides, it acts in the blood-current directly as an antiseptic. Prosorowsky (Meditzinskoje Obozrenije, No. 1, '98).

SOMATOSE.

Fifteen cases observed in which somatose was given to mothers whose supply of milk was failing. The result was good in those cases in which the somatose improved the appetite and general condition. Somatose, however, has no specific influence on the secretion of the mammary gland. Many women who are anxious to suckle their children are driven reluctantly to use the bottle, and in cases of this kind somatose is probably well worth trying. Joachim (Centralb. f. innere Med., March 12, '98).

SORE NIPPLES, THE PREVENTION OF.

The following method has proved unusually successful in the prevention of sore nipples:—

R Lanolin (Liebreich), 1 ounce.

Dispense in glass or porcelain screw-cap jar.

Sig.: For external use every night.

The patient is instructed to begin its use from four to six weeks before the expected date of confinement and continue until delivery. Every night at bedtime a small portion of lanolin is thoroughly worked into each nipple with the thumb and fingers, special pains being

taken to rub it well into any folds or crevices, especially in the case of depressed and sunken nipples.

In the morning it should be removed by a soft nail-brush, which is well soaked. The nipple should be brushed with lukewarm water and any mild, pure soap (preferably a white soap), giving it a thorough lathering for three or four minutes. It should afterward be rinsed with fresh water and dried as after ordinary bathing. All these agencies combined develop the cuticle, render it firm, elastic, and resisting, and produce a useful nipple, which may be almost guaranteed against subsequent abrasions and tenderness. J. Milton Mabbott (*New York Medical Journal*, Sept. 10, '98).

STARCH, SALIVARY DIGESTION OF.

Porridge with milk forms a more digestible compound, as far as amylolysis by saliva is concerned, than other combinations of oatmeal. The more dense, the less broken down, or the firmer the jelly in which the starchy food is when undergoing salivary digestion, the less rapid and extensive is the proteolysis. Some forms of starchy food undergo a certain amount of digestion by saliva; but at a certain stage this process stops, and, no matter how long they may be exposed to the action of ptyalin, no further amylolysis by saliva takes place (we must remember, however, the restraining effects which the products of digestion exert). Amylaceous substances are more easily acted on by saliva when thoroughly moist than when more or less dry. Bread in a light and spongy condition is more rapidly acted upon by saliva than when less spongy. Such bread, however, does not ultimately undergo any more complete digestion than does ordinary bread. Milk has a retarding influence on the salivary digestion of starch in bread,

while broth has little or no effect. Tea has a markedly inhibitory influence on amylolysis by saliva. Coffee has this property also to a less extent. Cocoa has hardly any restraining effect. Beer promotes the salivary digestion of starch. Alcohol, even in dilute solution, retards salivary digestion of starch, but the action is much less marked than in the case of infusions of tea. Wines have a very marked inhibitory influence on the digestion of starch by saliva, and this is almost wholly due to their acidity. W. G. Aitchison Robertson (*Jour. of Anat. and Phys.*, July, '98).

TYPHO-MALARIAL FEVER.

In India the combination of malarial infection with that of typhoid fever serves to obscure the early symptoms of typhoid, giving, in the majority of cases, a violent and sudden access, instead of the usual gradual step-like rise of the first three or four days. In such cases the plasmodium malariae may be detected in the blood during the first week, and again during the first week of convalescence (Laveran), introducing the intermittent character so often seen at those stages of typhoid fever in India. Beyond giving rise to this increased difficulty in diagnosis and delaying convalescence, the moderate presence of the malarial habit has no appreciable effect on the progress of an ordinary case of typical typhoid. A. Crombie (*British Medical Journal*, Sept. 24, '98).

TYPHOID FEVER.

Treatment.—Calomel is an exceedingly valuable drug in the treatment of typhoid fever, being directed against the seat of the disease. Its use is free from danger. It is best employed in small doses given hourly without interruption until all the symptoms of the disease are

entirely controlled or until sore gums warn us to discontinue it. Large purgative doses may be indicated early in the disease to absorb or modify it, or at any time to dissipate symptoms of profound toxæmia. The course of typhoid fever is greatly shortened by the systematic administration of calomel, and its graver features avoided. Henry W. Bettman (Nashville Jour. Med. and Surg., July, '98).

UTERINE CARCINOMA.

Treatment.—In uterine carcinoma the necrotic *débris* should be thoroughly removed by curetting under anæsthesia till firm tissue is reached. Hæmorrhage from arterial twigs is often checked by the use of the Paquelin cautery. Free irrigation with very hot water is then used to check the oozing, if necessary. It is desirable to have the seat of the operation as dry as possible before using carbide of calcium. Into the cavity which extends to a greater or less extent up into the body of the uterus, a piece of carbide, about the size of the last phalanx of one's thumb, is placed. At once acetylene-gas is evolved, which quickly fills the entire cavity with a froth, like soap-bubbles. The cavity is at once packed firmly with iodoform gauze. The vagina is packed full of the same material down to its ostium. The patient is then put to bed for three days, when the gauze and the carbide-remains are removed and a new piece is used. The carbide-remains are naught but a grayish clay covering the cavity in the body of the uterus. It can be sponged and irrigated away in a few minutes, dried by sponging, and prepared thus for another application of the carbide. Upon depositing the carbide in the wound, the same bubbling of gas is witnessed as before. Gauze packing is then

used and the patient is permitted to remain three days until the next dressing. After a series of such applications of the dressings the ragged, necrotic-faced ulcer is converted into a simple, clean ulcer. It presents a base covered, to all appearances, with pinkish-red granulations, destitute of the gray color which characterized its appearance after the second or third dressings. After a few treatments the edges of the cavity begin to draw in, and the area of the crater is diminished. Its entire appearance impresses one with the idea that it has taken on an entirely healthy character. Further persistence in the use is followed by progressive contraction, till finally the cavity is entirely obliterated. There is a puckering of the vault of the vagina about the small uterine os that remains, the whole field being covered by healthy, pink, mucous membrane. J. H. Etheridge (Jour. Amer. Med. Assoc., July 9, '98).

VARICOSE VEINS.

Trendelenburg's operation is the ideal method of treating varicose veins of the lower extremity associated with extensive ulceration. Trendelenburg found by experiments that the veins of the leg, after they had been temporarily emptied by elevation of the limb and compression of the trunk of the saphenous vein, are refilled slowly by the return blood coming from the arteries, and instantly by a blood wave coming from above downward. The conclusion that the veins in the leg are distended by great central pressure led this surgeon to advise ligation of the saphenous vein at two points and excision of the vessel between the ligatures. An incision about 4 inches in length is made over the saphenous trunk, beginning just above the union of the lower with the middle third of the

thigh. The vein, having been exposed, is carefully freed with a blunt dissector, and all branches going off from the vessel are ligatured. A ligature is then placed on the venous trunk at the upper

and another at the lower angle of the skin incision, and the portion of vein between these two ligatures is cut away with scissors. Cumston (*Annals of Surgery*, May, '98).

New Books Received.

The editor begs to acknowledge, with thanks, the receipt of the following books:—

Conservative Gynecology and Electro-therapeutics. A Practical Treatise on the Diseases of Women and Their Treatment by Electricity. Third Edition, Revised, Rewritten, and Greatly Enlarged. By G. Betton Massey, M.D., Physician to the Gynecic Department of Howard Hospital, Philadelphia; Late Electro-therapeutist to the Infirmary for Nervous Diseases, Philadelphia; Fellow and ex-President of the American Electro-therapeutic Association, of the Société Française d'Electrotherapie, of the American Medical Association, etc. Illustrated with Twelve Full-Page Original Chromolithographic Plates in Twelve Colors, Numerous Full-Page Original Half-tone Plates of Photographs from Nature, and many other Engravings in the Text. Royal Octavo. 400 pages. The F. A. Davis Co., Publishers, 1914-16 Cherry St., Philadelphia; 117 W. Forty-second St., New York City; 9 Lakeside Building, 218-220 S. Clark St., Chicago, Ill.—Special Report of the Beet-sugar Industry in the United States. Department of Agriculture, Washington, D. C., 1898.—Year-book of the United States Department of Agriculture, 1897.—Diphtheria und Diphtheritischer Croup. Von Dr. Adolf Baginsky. Alfred Holder, Wien, Publisher, 1898.

Monographs Received.

The editor begs to acknowledge, with thanks, the following list of monographs:—

Equilibration and its Relation to Vertigo. By Frank K. Hallock, A.M., M.D., Cromwell, Conn., 1898.—Injuries of the Newborn in Cross, Complex, and Breech Presentations. By Edwin Rosenthal, M.D., Philadelphia, 1898.—The Dosage of Diphtheria Antitoxin and Its Method of Using. By Edwin Rosenthal, M.D., Philadelphia, 1898.—Diseases of Tropical Climates: Their Prevention, Diagnosis, and Treatment. By T. S. Dabney, M.D., New Orleans, 1898.—Intubation in Diphtheria. By W. K. Simpson, M.D., New York, 1898.—An Improved Form of Stereoscope. By Charles A. Oliver, A.M., M.D., Philadelphia, 1898.—Progressive Loss of Brain-weight in Dementia. By Warren L. Babcock, M.D., Ogdensburg, N. Y., 1898.—New Forceps for Intestinal Anastomosis. By Ernest Laplace, M.D., LL.D., Philadelphia, 1898.—Report of a Case of Acute Double Hydrocele, due to Secondary Syphilis. By Howard Paxton Collings, B.S., M.D., Hot Springs, Ark., 1897.—The Periodical Cicada; an Account of Cicada Septendecim, its Natural Enemies and the Means of Preventing its Injury, together with a Summary of the Distribution of the Different Broods. By C. L. Marlett, M.S., U. S. Department of Agriculture, Washington, D. C., 1898.—Contributo Alla Statistica, Dell' ascesso Peritracheo-Laringeo Nei Bambini, Descritto Dal Massei. By Dott. Francesco Egidi, Rome, 1898.—Contributo Clinico ed Anatomo-pathologico, allo Studio

Dell' emicorea Post-emiplegica E dei Rammollimenti del Ponte di Varolio. By Dott. Alfredo Rubino, Naples.—Sull' Epilessia Sifilitica Secondaria. By Dott. Alfredo Rubino, Naples, 1898.—Neurotic Eczema. By L. Duncan Bulkley, A.M., M.D., New York, 1898.—Errors in Death-returns from Malarial and Typho-malarial Fevers; Typhoid Fever the Prevailing Fever of Washington. By William W. Johnston, M.D., Washington, D. C., 1898.—The Influence of School-life upon the Health of Children. By William W. Johnston, M.D., Washington, D. C., 1897.

EDITORIAL STAFF


OF

Sajous's Annual and Analytical Cyclopædia of Practical Medicine.

ASSOCIATE EDITORS.

(List Revised August 1, 1898.)

J. GEORGE ADAMI, M.D., MONTREAL, P. Q.	S. G. GANT, M.D., KANSAS CITY, MO.	WILLIAM OSLER, M.D., BALTIMORE, MD.
LEWIS H. ADLER, M.D., PHILADELPHIA.	J. McFADDEN GASTON, M.D., ATLANTA, GA.	F. A. PACKARD, M.D., PHILADELPHIA.
JAMES M. ANDERS, M.D., LL.D., PHILADELPHIA.	J. E. GRAHAM, M.D., TORONTO, ONT.	LEWIS S. PILCHER, M.D., BROOKLYN, N. Y.
G. APOSTOLI, M.D., PARIS, FRANCE.	JULES GRAND, M.D., PARIS, FRANCE.	WILLIAM CAMPBELL POSEY, M.D., PHILADELPHIA.
A. D. BLACKADER, M.D., MONTREAL, P. Q.	EGBERT H. GRANDIN, M.D., NEW YORK CITY.	W. B. PRITCHARD, M.D., NEW YORK CITY.
E. D. BONDURANT, M.D., MOBILE, ALA.	LANDON CARTER GRAY, M.D., NEW YORK CITY.	JAMES J. PUTNAM, M.D., BOSTON.
DAVID BOVAIRD, M.D., NEW YORK CITY.	J. P. CROZER GRIFFITH, M.D., PHILADELPHIA.	GEORGE H. ROHÉ, M.D., SYKESVILLE, MD.
L. BROCC, M.D., PARIS, FRANCE.	A. GOUGUENHEIM, M.D., PARIS, FRANCE.	ALFRED RUBINO, M.D., NAPLES, ITALY.
WILLIAM BROWNING, M.D., BROOKLYN, N. Y.	C. M. HAY, M.D., PHILADELPHIA.	LEWIS A. SAYRE, M.D., NEW YORK CITY.
WILLIAM T. BULL, M.D., NEW YORK CITY.	FREDERICK P. HENRY, M.D., PHILADELPHIA.	REGINALD H. SAYRE, M.D., NEW YORK CITY.
CHARLES W. BURR, M.D., PHILADELPHIA.	EDWARD JACKSON, M.D., DENVER, COL.	SOLOMON SOLIS-COHEN, M.D., PHILADELPHIA.
DUDLEY W. BUXTON, M.D., M.R.C.P. LONDON, ENGLAND.	NORMAN KERR, M.D., F.L.S., LONDON, ENGLAND.	H. W. STELWAGON, M.D., PHILADELPHIA.
HENRY T. BYFORD, M.D., CHICAGO, ILL.	EDWARD L. KEYES, JR., M.D., NEW YORK CITY.	D. D. STEWART, M.D., PHILADELPHIA.
J. ABBOTT CANTRELL, M.D., PHILADELPHIA.	H. KRAUSE, M.D., BERLIN, GERMANY.	LEWIS A. STIMSON, M.D., NEW YORK CITY.
WILLIAM B. COLEY, M.D., NEW YORK CITY.	E. LANDOLT, M.D., PARIS, FRANCE.	G. ARCHIE STOCKWELL, M.D., NEW YORK CITY.
P. S. CONNER, M.D., LL.D., CINCINNATI, OHIO.	ERNEST LAPLACE, M.D., LL.D., PHILADELPHIA.	B. J. STOKVIS, M.D., AMSTERDAM, HOLLAND.
FLOYD M. CRANDALL, M.D., NEW YORK CITY.	R. LÉPINE, M.D., LYONS, FRANCE.	LOUIS McLANE TIFFANY, M.D., BALTIMORE, MD.
ANDREW F. CURRIER, M.D., NEW YORK CITY.	F. LEVISON, M.D., COPENHAGEN, DENMARK.	CHARLES S. TURNBULL, M.D., PHILADELPHIA.
JUDSON DALAND, M.D., PHILADELPHIA.	A. LUTAUD, M.D., PARIS, FRANCE.	HERMAN F. VICKERY, M.D., BOSTON, MASS.
N. S. DAVIS, M.D., CHICAGO, ILL.	F. MASSEI, M.D., NAPLES, ITALY.	RIDGELY B. WARFIELD, M.D., BALTIMORE, MD.
F. EKLUND, M.D., STOCKHOLM, SWEDEN.	E. E. MONTGOMERY, M.D., PHILADELPHIA.	F. E. WAXHAM, M.D., DENVER, COL.
AUGUSTUS A. ESHNER, M.D., PHILADELPHIA.	JULES MOREL, M.D., GHENT, BELGIUM.	J. WILLIAM WHITE, M.D., PHILADELPHIA.
J. T. ESKRIDGE, M.D., DENVER, COL.	HOLGER MYGIND, M.D., COPENHAGEN, DENMARK.	W. NORTON WHITNEY, M.D., TOKIO, JAPAN.
CHRISTIAN FENGER, M.D., CHICAGO, ILL.	W. P. NORTHRUP, M.D., NEW YORK CITY.	JAMES C. WILSON, M.D., PHILADELPHIA.
SIMON FLEXNER, M.D., BALTIMORE, MD.	H. OBERSTEINER, M.D., VIENNA, AUSTRIA.	C. SUMNER WITHERSTONE, M.D., PHILADELPHIA.
LEONARD FREEMAN, M.D., DENVER, COL.	CHARLES A. OLIVER, M.D., PHILADELPHIA.	WALTER WYMAN, M.D., WASHINGTON, D. C.



FORMULA:

R Sumbul.
 Iron.
 Cinchona.
 Acid Phosp.
 Aromatics. } Qs.
 Sherry Wine.

TONO SUMBUL CORDIAL

(W. R. WARNER & CO.)

Comprises a combination of tonics in active form, which, taken under the supervision of the physician, will cause patients to gain in strength and weight.

As will be seen, it does not contain Coca or any ingredient which might induce a drug habit, but is a superior tonic, used to advantage and discontinued with no after effects.

It is palatable, and especially serviceable where the stomach of the invalid rebels at a nauseating preparation.

SEE THAT YOU GET THE ORIGINAL.

SUPERIOR TO PEPSIN OF THE HOG.

INGLUVIN

A Powder—prescribed in the same manner, doses and combinations as Pepsin.

A Specific for Vomiting in Pregnancy.

PREPARED ONLY BY

W. R. WARNER & CO.

PHILADELPHIA.

NEW YORK.

CHICAGO.

ELIXIR CHLORALAMID

AN EFFICIENT AND SAFE HYPNOTIC.

Each Tablespoonful contains 15 grains Chloralamid.

The large number of physicians who have recorded their experience with Chloralamid agree that it is a potent hypnotic free from any depressing action on the heart. Elixir Chloralamid presents the remedy in most suitable form for administration. In doses from 15 to 45 grains it induces sound sleep lasting from 5 to 8 hours.

Chloralamid is official in the German Pharmacopœia and is endorsed by Prof. Chas. L. Dana, Prof. John V. Shoemaker, Dr. H. C. Wood, Dr. David Cerna, Dr. John Aulde, Prof. Chas. H. Steele, Prof. R. W. Wilcox, and others.

LEHN & FINK, Manufacturing Chemists, NEW YORK.

In writing to advertisers mention Monthly Cyclopædia

NOTES AND COMMENTS.

THE F. A. DAVIS COMPANY, Publishers.

GEO. B. CREVELING, Business Manager.

Hawaii and the Philippines.—Send four cents (in stamps) for an illustrated booklet issued by the Chicago, Milwaukee & St. Paul Railway, the direct route across the American Continent to the new Transpacific possessions of the United States. Full of latest reliable information and valuable for reference. Can be used as a text-book in school. Address, John R. Pott, District Passenger Agent, Williamsport, Pa.

An Extraordinary Offer.—In order to thoroughly introduce "Unguentine" to the medical profession, the Norwich Pharmacal Company, Norwich, N. Y., will send one of their collapsible tubes (see their advertisement, page 9) free of expense, to all of our subscribers who will mention this publication. We consider this an exceedingly liberal offer, and as "Unguentine" is favorably known, we trust that our readers will take advantage of this offer.

New Uses of Antikamnia.—A 5-grain-antikamnia tablet taken before one starts upon a pleasure-trip will entirely prevent the demoralizing headache which frequently mars the enjoyment of such an occasion. The nervous headache and irritable condition of the overworked businessman is prevented by the timely use of a 10-grain dose.

The Pioneer Limited is the name of the only perfect train in the world, now running every night between Chicago, St. Paul, and Minneapolis *via* the Chicago, Milwaukee & St. Paul Railway:

the pioneer road of the West in adopting all improved facilities for the safety and enjoyment of passengers. An illustrated pamphlet, showing views of beautiful scenery along the route of the Pioneer Limited, will be sent free to any person upon receipt of a two-cent postage stamp. Address Geo. H. Heafford, General Passenger Agent, Chicago, Ill:

Where Noah Kept His Bees.—Dr. James K. Hosmer, while recently visiting Boston, had occasion to visit the new public library. As he went up the steps he met Edward Everett Hale, who asked the doctor's errand.

"To consult the archives," was the reply.

"By-the-way, Hosmer," said Doctor Hale, "do you know where Noah kept his bees?"

"No," answered Hosmer.

"In the ark hives," said the venerable preacher as he passed out of earshot.—"Summer Piazza Stories" in the August Ladies' Home Journal.

An Editor's Daughter's Dolls.—The editor of Harper's Magazine, Mr. Henry M. Alden, has a daughter who has made a collection of over 100 dolls, each doll representing a different nation and being made in that country. The collection is considered to be the most unique and finest of its kind in existence. Miss Alden is now to show the world her dolls, and explain them, and in the next issue of The Ladies' Home Journal she will show pictures of the first twenty. In following issues she will show the others.

In writing to advertisers mention Monthly Cyclopædia

Rx: Ung't. Resinol \mathfrak{z} i—iii.

THOROUGHLY EFFICIENT AS AN
ANTIPRURITIC, ANTISEPTIC,
LOCAL ANTIPHLOGISTIC & SKIN NUTRIENT

And is now being successfully
used and prescribed by over

40,000 PHYSICIANS in the United States
and Canada

For Eczema, Erysipelas, Pruritus and all Itching, and
Inflammatory Skin Diseases.

Rx: Elix. Cascanata \mathfrak{z} iv—x.

LAXATIVE, ALTERATIVE, ANTACID AND TONIC.

Stimulates the elimination of effete matter and restores the
normal alkalinity of the blood.

A specific for habitual constipation and the digestive derangements resulting therefrom.

RESINOL CHEMICAL CO., = Baltimore, Md.

When a Physician

Perry Patetic — They
say a man enjoys restin'
a whole lot better after a
good hard day's work.

Wayworn Watson —
Well, fer all I know, it
may be so; but I ain't
'round tryin' dangerous
experiments.

has learned by experience that LACTO-
PEPTINE is almost a specific in the
treatment of digestive disorders it's a
"dangerous experiment" to allow the
substitution of any of its imitations
when LACTOPEPTINE is prescribed.
Dangerous to the patient, whose wel-
fare may depend upon its prompt effects;
dangerous to the physician, whose repu-
tation may suffer by failure to afford
relief; dangerous to the 'pharmacist
should either patient or physician learn
of the imposition practiced. Avoid these
dangers, Doctor, by insisting that the
genuine LACTOPEPTINE is dis-
pensed when prescribed. It has never
been successfully imitated.

THE NEW YORK PHARMACAL ASSOCIATION,
YONKERS, N. Y.

Samples upon request.

In writing to advertisers mention Monthly Cyclopædia

Notes and Comments (continued).

Grape-juice for Patriots.—The Welch Grape-juice Company have given away over one hundred cases of grape-juice to the government, at Santiago and Montauk Point, and to hospitals where sick soldiers are being nursed to health.

Oak-Grove Hospital.—One of the happiest and most beneficent reforms of modern times relates to the treatment of the insane. In addition to those deprived of the use of their reason there are countless sufferers from nervous maladies who are entirely unfit for the struggle for existence. Perhaps a more pitiable fate than actual insanity is to be an imbecile in a slight degree, but just enough to be rendered incompetent for the serious business of life.

For the humane treatment of these unfortunates by hygienic, medicinal, and moral means the Oak-Grove Hospital, situated at Flint, Michigan, offers many advantages. The hospital is located in the midst of an ample and well-wooded domain. The apartments are well furnished, the attendance is excellent, and every arrangement has been made for the use of baths, massage, electricity, etc. The institution is also provided with gymnasium, billiard-room, bowling-alley, etc., for the use of those to whose cases and condition they may be properly adapted.

A New Vegetable Alterative Compound.

—The Resinol Chemical Company, of Baltimore, have prepared a scientific combination called Cascanetta, which contains the active principles of cascara sagrada, gentian, rhubarb, trifolium, and rumex. The Elixir Cascanetta is recommended as highly serviceable in cutaneous disorders due to morbid conditions of the alimentary tract. Disorders result-

ing from congestion of the pelvic viscera are also alleviated by this preparation.


Axion Elastic Truss.—The axion truss is made with a superior-grade elastic-silk webbing, manufactured exclusively for use in this truss. The elastic is strong, durable, and soft. The new Axion pad gives universal satisfaction, being perfectly adjustable in every way and adapted to every variety of hernia.

Antikamnia in Acute Prostatitis.—In the course of a paper read in the Section of Surgery and Anatomy of the American Medical Association by Dr. Liston Homer Montgomery, of Chicago, antikamnia was recommended as a reliable anodyne remedy to control pain and spasm of the neck of the bladder.

Bensolyptus.—Bensolyptus is the name given by Schieffelin & Co., of New York, to a preparation containing efficient antiseptic substances as well as alkaline salts. This preparation is highly esteemed for its beneficial influence in diseases of the mucous membrane used both internally and by the mouth.

Maltzyme.—Maltzyme is an improved essence of malt, and is very useful in the treatment of starchy indigestion. It can be administered alone or, where such combinations are indicated, with cod-liver-oil, cascara sagrada, and the hypophosphites.

Baird's Steam-fomenter.—This is an apparatus which should be in every household. It generates a continuous current, and its construction is so simple that it can be managed by any one. An apparatus of this kind is useful in every condition where heat is required, and



ABDOMINAL SUPPORTING BELTS for every condition requiring abdominal support. Our Belts are made to measurements, and they fit; consequently, they are efficient. Price lists and order blanks sent on request.

SPECIALTIES.—Pomeroy Frame Truss, "Master" Elastic Stockings.

POMEROY COMPANY.

19 W. Union Square, New York City.

JUST READY.

New, Revised Edition of an exceedingly useful
... book ...

A Primer of Psychology and Mental Disease

FOR USE IN TRAINING-SCHOOLS FOR ATTENDANTS AND NURSES
AND IN MEDICAL CLASSES.

By **C. B. BURR, M.D.,**

Medical Director of Oak Grove Hospital for Nervous and Mental Diseases, Flint, Mich. Formerly Medical Superintendent of the Eastern Michigan Asylum. Member of the American Medico-Psychological Association, etc.

Second Edition. Thoroughly Revised and Illustrated. 116 pages, 5¼ x 8 inches. Neatly Bound in Extra Cloth. Price, \$1.00, Net, POST-PAID.

THE F. A. DAVIS COMPANY, PUBLISHERS, 1914 AND 1916 CHERRY STREET, PHILADELPHIA.

ESPERANZA (DRY).



HYGEIA (SWEET).

Unfermented Concord Grape Juice.

FROM CHOICE, SELECTED GRAPES. NO
ALCOHOL. NO ANTISEPTICS.

Nature's Best Drink and Tonic.

FOR THE SICK. FOR THE WELL.

Prepared by

SNOW GRAPE JUICE CO.,

PENN YAN, N. Y.

SEND 30 CENTS IN STAMPS FOR PINT BOTTLE.

In writing to advertisers mention Monthly Cyclopædia

Notes and Comments (continued).

may be employed as a vapor-bath or by inhalation, according to the circumstances of each particular case. Baird's steam-fomenter can be obtained from Baird Brothers Company, of Chicago.

Chionia for the Liver.—Those who possess what is called the bilious temperament and likewise all who do not have sufficient exercise to render the circulation vigorous and stimulate the liver will be benefited by chionia, which has a special selective action upon the hepatic cells. Chionia is not a purgative, but it brings about a natural regularity of the bowels. The dose of this preparation is 1 to 2 fluidrachms three times a day, and it is made by the Peacock Chemical Company, of St. Louis.

Uterotonic (N. B. & Co.).—Uterotonic is an efficient remedy in all conditions of the female generative apparatus which are amenable to the action of internal remedies. In cases which require surgical intervention uterotonic may be properly used as an adjuvant which has a tonic and beneficial influence upon the organs of the pelvis.

Free Treatise on Electrotherapy.—The reputation of the McIntosh Battery and Optical Co., of Chicago, is of the

highest order. Our readers will consequently be glad to learn of the generous offer made by such a reliable company. They offer to send post-paid and free of charge, to anyone who writes asking for it, a treatise of three hundred pages and four hundred and fifty illustrations. The book gives the technique of the various operations performed by means of electricity.

Imperial Granum.—This preparation, so favorably known to the medical profession, made an extremely-attractive display at the Denver meeting of the American Medical Association. The imperial granum is an excellent article, and wins friends wherever it goes.

Xeroform in Diseases of the Skin.—Dr. Ehrmann, of Vienna, has used xeroform for more than a year, and has treated 178 patients with it externally and 45 patients internally; as 13 received the remedy both externally and internally the entire number is equivalent to 210. His cases included superficial diseases, suppurations and necroses, and clean operative wounds. All the operative wounds healed by first intention. Secretion diminished rapidly in varicose ulcers. Xeroform was equally beneficial in moist eczema. Internally, the author gave xeroform in several diseases of the



The Trained Nurse Knows
Platt's Chlorides.
The True Disinfectant.

Platt's Chlorides, ***The True Disinfectant.***

An odorless, colorless liquid; powerful, safe, and cheap; indorsed by over 23,000 physicians; sold everywhere in quart bottles only. Prepared only by

HENRY B. PLATT,

Platt Street, New York.

In writing to advertisers mention Monthly Cyclopædia

Notes and Comments (continued).

skin notoriously accompanied by increased decomposition of the intestinal contents. Xeroform has no unpleasant odor or taste and caused no eructations. It was given in 8-grain doses three or four times a day.

Pepto-mangan in Surgical Convalescence.—Dr. Stuart McGuire, of Richmond, has tested the action of Gude's Pepto-mangan in twenty major cases on which he had operated. The period during which the preparation was used in these cases varied from ten to sixty days.

The results were exceedingly satisfactory. Dr. McGuire says that when it is remembered that the patients were all confined to bed, that they were recovering from the effects of serious surgical operations, and that they were subjected to the depressing influence of hospital-life, the average increase of red blood-corpuscles is remarkable. In all the cases the number of red blood-cells was accurately determined by count.

Kryofine in Typhoid Fever and Insomnia.—Kryofine is a valuable remedy in typhoid fever because it reduces the

VACCINE

PURE AND RELIABLE ANIMAL VACCINE LYMPH.
Fresh Daily. Send for Illustrated Pamphlet.
10 IVORY POINTS, Double Charged, \$1.00. 10 CAPIL-
LARY TUBES, \$1.00. Orders by Mail or Tele-
graph Promptly Dispatched.

VIRUS.

NEW ENGLAND VACCINE CO., Chelsea Station, BOSTON, MASS.

WM. C. CUTLER, M.D.

CHAS. N. CUTLER, M.D.

AS AN ANODYNE AND HYPNOTIC

...KRYOFINE...

(Methoxacet-p-phenetidin)

IS POWERFUL AND WITHOUT ILL EFFECTS.

DR. GEORGE F. BUTLER, Professor of Materia Medica and Clinical Medicine, College of Physicians and Surgeons, Chicago; Attending Physician Cook County Hospital, etc., in the course of an article in the Chicago Clinic, says:

"With the exception of morphine, no drug possesses so positive, prompt and efficient an analgesic property as Kryofine; indeed there are painful disorders—such as migraine, and particularly the pains of locomotor ataxia and certain spinal diseases—where Kryofine seems nearly as efficient as morphine, and attended with less unpleasant sequelæ."

In the services of Drs. J. Rudisch, A. Meyer and A. G. Gerster, Mt. Sinai Hospital, New York, it was found that:

"As a hypnotic in insomnia, unaccompanied by severe pain, it has proved of undoubted value."

"In a case of acute suppurative arthritis, general sepsis and pericarditis, the drug did not affect the pulse. When the child was restless at night, 2½ grains of Kryofine produced a quiet sleep, lasting several hours."—Drs. HAAS and MORRISON, New York Med. Journal.

Dose:—4-7½ grs., tablets or crystalline powder.

Literature and Samples sent on request.

C. BISCHOFF & CO., NEW YORK.

In writing to advertisers mention Monthly Cyclopædia

Notes and Comments (continued).

temperature promptly and keeps it down for a prolonged period. It increases arterial tension, has no deleterious action on the kidneys, acts favorably in patients of any age, has a marked hypnotic influence, and lessens or relieves delirium, stupor, or subsultus. In insomnia, whether dependent or not upon pain, the members of the staff of Mt. Sinai Hospital reported very satisfactory results. Led by this report Dr. I. J. Higgins made use of kryofine in an extremely obstinate case of insomnia. He administered rather more than 6 grains every night and found it effectual. For the second week it was directed to be taken every other night, which proved equally serviceable. The author reports this case with the view of interesting other observers in the clinical investigation of the drug.

"A New Created World."—If you leave home for the fall, go where you will find almost literally a new world. Such a spot is Yellowstone Park. There is no place like it in the world. You will be astonished at what you see and learn there. A month's outing in those mountains 7000 feet above the sea-level, in an atmosphere pure and invigorating, would add years to your life. There are four modern hotels, good roads—maintained by the United States Government—for riding, driving or cycling; riding-horses and pack-trains. Geysers, canyons, mud-springs, lakes, cataracts, and cascades in large number are crowded into a space 54 by 62 miles in size. If you can afford it, you wrong yourself by not going there, especially if you are a patriotic American. If you cannot afford it, then go to beautiful Leech Lake, in Minnesota, a new and popular resort. New hotels, fine fishing

ELIXIR MALTOPEPSINE (TILDEN'S)

DIGESTS ALL KINDS OF FOOD.

Is especially useful in **CHOLERA INFANTUM** and all forms of **SUMMER COMPLAINT WITH CHILDREN**. Promotes digestion by aiding and perfecting the process. Gives immediate relief in **DYSPEPSIA**, and aids in curing it. Send for samples and literature.

RESPIRAZONE (TILDEN'S)

Indispensable in the treatment of **ACUTE** and **CHRONIC ASTHMA**, **HAY FEVER**, and **CROUP**. Physicians everywhere are prescribing this preparation with unvarying results. It is not a new remedy, but a correct combination of old remedies, and the profession can prescribe it feeling confident of results. Doctor, if you have not used **RESPIRAZONE (TILDEN'S)** send at once for samples and literature.

MANUFACTURED ONLY BY

THE TILDEN COMPANY,

MANUFACTURING PHARMACISTS,

NEW LEBANON, N. Y.

ST. LOUIS, MO.

In writing to advertisers mention Monthly Cyclopædia

Notes and Comments (continued).

and hunting, and good boating at reasonable rates. Send Charles S. Fee, of Northern Pacific Railway, St. Paul, Minn., six cents for Wonderland, '98, that describes by pen and picture both resorts. Mention where advertisement was seen.

Profits of Railways.—The Report of the Interstate Commerce Commission shows that last year the railways of the United States carried over 13,000,000,000 passengers 1 mile. They also carried 95,000,000,000 tons of freight 1 mile. The total amount paid in dividends on stock was \$87,603,371—call it \$88,000,000. Of the total earnings of the railways about 70 per cent. came from the freight service, and 30 per cent. from passenger-service. Let us assume, then, that of the \$88,000,000 paid in dividends, 70 per cent., or \$61,600,000 was profit

on freight-service and \$26,400,000 was profit on passenger-service. Let us drop fractions and call it \$62,000,000 from freight and \$26,000,000 from passengers. By dividing the passenger-profit into the number of passengers carried (13,000,000,000) we find that railways had to carry a passenger 500 miles in order to earn one dollar of profit—or 5 miles to earn one cent. Their average profit, therefore, was less than two-tenths of one cent for carrying a passenger (and his baggage) 1 mile.

By dividing the freight-profit into the freight mileage (95,000,000,000) we find that the railways had to carry one ton of freight 1,530 miles in order to earn one dollar, or over 15 miles to earn one cent. The average profit therefore was less than one-fifteenth of a cent for carrying a ton of freight (besides loading and unloading it) 1 mile.

White Rock Ozonate Lithia Water

PLEASES PARCHED PALATES PERFECTLY.



People drink it because they like it; hence the uniform success reported by thousands of physicians who prescribe it.

Bottled Only by

WHITE ROCK MINERAL SPRINGS CO., Waukesha, Wis.

In writing to advertisers mention Monthly Cyclopædia

Notes and Comments (continued).

Wampole's Liquid Wheat.—This preparation has been introduced by Henry K. Wampole & Co., of Philadelphia, as a nutrient tonic for use in neurasthenia, lactation, indigestion, to maintain nutrition in cholera infantum, to promote development of the teeth, etc. Being palatable, it is readily taken by children.

It is described as a concentrated, predigested preparation of the whole wheat-grain, and contains over 7 per cent. of the nutrient vitalized hypophosphites and oleonitrogenous hypophosphites obtained solely from wheat, principally from the germ and cortical portion,—the parts usually rejected in making wheat-flour. It is rich in the flesh-producing gluten and albumin, contains over 4 per cent. of oil of wheat, and the wheat-starch is converted into maltose,—a direct force- and fat-making product. Especially valuable for nervous, backward children.

Each teaspoonful represents 1 ounce of the entire wheat-grain, in perfect solution, and $\frac{1}{8}$ grain U. S. P. pepsin.

The dose for adults is 1 teaspoonful, gradually increasing to 1 tablespoonful, three or more times daily, with or after meals. For children 3 to 5 years old, commence with $\frac{1}{2}$ teaspoonful, gradually increasing to 1 teaspoonful.

The Safety of Railway Travel.—Last

year on American railways one passenger was killed in accidents out of every 2,827,474 passengers carried. That is to say, that you can take a train 2,827,474 times before, on the law of averages, your turn comes to be killed. You will have to travel 72,093,963 miles on the cars before that turn comes, and 4,541,945 miles before you are injured. If you travel 20 miles every day for 300

days in the year, you can keep on at it for 758 years before your turn comes to be hurt. If there had been railways when our Saviour was born and you had begun to travel on the first day of the year A.D. 1, and had traveled 100 miles in every day of every month of every year since then, you would still have (in this year 1898) nearly three million miles yet to travel before your turn came to be killed.

Fever in Infectious Diseases.—To reduce

the fever in attacks of pneumonia, typhoid, gastric fever, scarlet fever, measles, etc., the following will be found to act promptly, with grateful calmative effect:—

For adults:

℞ Lactaphenin, 1 $\frac{1}{4}$ drachms.

Divide into 10 doses (tablets, powders, or capsules).

Sig.: Take 1 every three hours.

For children of 5 to 10 years:

℞ Lactophenin, 25 grains.

Sugar, 30 grains.

Divide into 10 powders and give 1 every three hours.

For children of 2 to 5 years:

℞ Lactophenin, 15 grains.

Sugar, 45 grains.

Divide into 10 powders and give 1 every three hours.



**DR. BECKER'S
COMPOUND
DIGEST.**

(Digestinla Composita. Dr. Becker.)

is prepared according to the formula of Dr. Becker, from principles obtained by his processes, from the digestive organs of clean animals and fowls.

It assists impaired and overworked digestive organs, allowing them to rest without lessening nutrition.

It is easily administered, and, being free from chemicals, does not interfere with the action of other medicines.

Sold only in bottles containing 550 grains powder form, or 100 five-grain Tablets, and may be ordered through any druggist. Samples sent by mail to physicians on application.

DR. BECKER COMPOUND DIGEST CO.,

107 Dearborn Street, CHICAGO, ILL.

In writing to advertisers mention Monthly Cyclopædia

THE SANITARIUM, BATTLE CREEK, MICH.

INCORPORATED 1867.



THE largest, most thoroughly equipped and one of the most favorably located in the United States. It is under strictly regular management. Eight physicians, well-trained and of large experience. A quiet home-like place, where "trained nurses," "rest-cure," "massage," "faradization," "galvanization," "static electrization," "Swedish movements," "dieting," "baths," "physical training," and all that pertains to modern rational medical treatment can be had in perfection at reasonable prices. Special attention given to the treatment of chronic disorders of the stomach, and diseases peculiar to women. A special hospital building (100 beds) for surgical cases with finest hospital facilities and appliances.

Large fan for winter and summer ventilation. Absolutely devoid of usual hospital odors. Delightful surroundings. Lakeside resort. Pleasure grounds, steamers, sailboats, etc. Trained nurses of either sex furnished at reasonable rates.

J. H. KELLOGG, M. D., Supt., Battle Creek, Mich.

R **NUT-GLUTEN BISCUIT:** Made from the purest wheat gluten and nut meal. The only food on the market to-day that diabetics may eat of freely, without danger. Prescribed and endorsed by the physicians of the Battle Creek Sanitarium. Samples free to physicians when card is enclosed with the request. Others may have them for two 2c stamps.

BATTLE CREEK SANITARIUM HEALTH FOOD CO., Battle Creek, Mich.



In writing to advertisers mention Monthly Cyclopædia

Artistic Home Decorations.

We can show you effects NEVER before thought of, and at moderate prices, too.

Why have your house decorated and painted by inferior workmen, when you can have it done by skilled workmen—by Artists—for the same price?

If you intend Decorating, if only one room, call or send to see what we are doing, and for whom.

TAPESTRY PAINTINGS.

2000 tapestry paintings to choose from. 38 artists employed, including gold medalists of the Paris Salon. Send 25 cents for compendium of 140 studies. When in the city be sure to call and see our exhibition.

WALL PAPERS.

New styles, designed by gold-medal artists. From 10 cents per roll up. Send 50 cents to prepay expressage on large sample books and drapery. Will include drapery samples in package. See our Antique Metallic, French, Pressed Silks, and Lida effects in special colors to match all kinds of wood-work, carpets, and draperies. Have 500 different wall-hangings with draperies specially made at our Broomhead Mills, Paterson, N. J., to match.

DECORATIONS.

Color-schemes—designs and estimates submitted free. Artists sent to all parts of the world to do every sort of decorating and painting. We are educating the country in color-harmony. Relief, stained glass, wall-paper, carpets, furniture, draperies, etc. Pupils taught.

DECORATIVE ADVICE.

Upon receipt of \$1 Mr. Douthitt will answer any question on interior decorations—color-harmony and harmony of form, harmony of wall-coverings, carpets, curtains, tiles, furniture, gas-fixtures, etc.

MANUAL OF ART DECORATIONS.

The art-book of the century. 200 royal quarto pages, filled with full-page colored illustrations of modern home interiors and tapestry studies. Price \$2.

If you want to be up in decoration send \$2 for this book; worth \$50.

SCHOOL.

Six 3-hour tapestry painting lessons, in studio, \$5. Complete written instructions by mail, \$1. Tapestry paintings rented; full-size drawings, paints, brushes, etc., supplied. Nowhere, Paris not excepted, are such advantages offered pupils. New catalogue of 125 studies, 25 cents.

Send \$1 for complete instructions in tapestry painting and compendium on 140 studies.

TAPESTRY MATERIALS.

We manufacture tapestry materials superior to foreign goods and half the price. Book of samples, 10 cents.

Send \$1.50 for 2 yards No. 6, 50-inch goods, just for a trial order; worth \$3. All kinds of draperies to match all sorts of wall papers, from 10 cents per yard up. **THIS IS OUR GREAT SPECIALTY.**

GOBLIN-PRINTED BURLAPS.

Over 100 new styles for wall-coverings, at 25 cents per yard, 35 inches wide, thus costing the same as wall-paper at \$1 per roll. 240 kinds of Japanese lida leather papers at \$2 per roll.

GOBLIN ART DRAPERY.

Grecian, Russian, Venetian, Brazilian, Roman, Rococo, Dresden, Festoon, College Stripe, Marie Antoinette, Indian, Calcutta, Bombay, Delft, Soudan.

In order that we may introduce this line of New Art Goods, we will send one yard each of 50 different kinds of our most choice Patterns for \$7.50.

JOHN F. DOUTHITT,

American Tapestry Decorative Co.,

286 Fifth Avenue, New York.

Near 30th Street.

In writing to advertisers mention Monthly Cyclopædia

Dietz Driving Lamp

only claims to be the very best Lamp in existence for driving after nightfall—'tis in the color, volume and penetration of the light it gives, and especially desirable because that light is wind and jolt-proof.




To its virtues in detail we devote a special Catalogue that will interest you—'tis mailed free, of course, as is also our Complete Catalogue.

Upon receipt of \$3.75 we will send you (freight prepaid) one of these Lamps with bracket for your carriage.

R. E. DIETZ CO.,
60 Laight Street, New York.

ESTABLISHED IN 1840.

First see "DIETZ" on any Lantern or Lamp
—then go ahead.



CLIMATE CURE

The **SALT RIVER**
VALLEY of
Arizona and the
various
HEALTH RESORTS
in **NEW MEXICO**

ARE UNRIVALED FOR THE RELIEF
OF CHRONIC LUNG AND THROAT DIS-
EASES. HERE ARE FOUND PURE DRY AIR,
EQUABLE TEMPERATURE AND CONSTANT SUNSHINE.
THE ITEMS OF ALTITUDE, TEMPERATURE, HUMIDITY, HOT SPRINGS,
SANATORIUMS, COST OF LIVING, MEDICAL ATTENDANCE, SOCIAL
ADVANTAGES, ETC., ARE CONCISELY TREATED IN DESCRIPTIVE
PAMPHLETS ISSUED BY THE SANTA FE ROUTE
PHYSICIANS ARE RESPECTFULLY ASKED TO PLACE THIS
LITERATURE IN THE HANDS OF INVALIDS.

ADDRESS **W.J. BLACK,**
G.P.A., A.T. & S.F. RY.,
TOPEKA, KANSAS.

OR **C.A. HIGGINS,**
A.G.P.A., CHICAGO.

NOW READY.

A Very Important Contribution to Laryngological Science.

Autoscopy of the Larynx and the Trachea.

(DIRECT EXAMINATION WITHOUT MIRROR.)

By **ALFRED KIRSTEIN, M.D.,** Berlin.

Authorized Translation (Altered, Enlarged, and Revised by the Author) by

MAX THORNER, A.M., M.D., Cincinnati, O.

Professor of Clinical Laryngology and Otology, Cincinnati College of Medicine and Surgery; Laryngologist and Aurist, Cincinnati Hospital, etc.

About 75 Pages, with Twelve Illustrations. Crown Octavo, bound in
Extra Cloth, price, 75 cents net.

THE F. A. DAVIS COMPANY, Publishers.

PHILADELPHIA: 1914 and 1916 Cherry St.

NEW YORK: 117 W. Forty-Second St.

CHICAGO: 9 Lakeside Building, 214-220 S. Clark St.

In writing to advertisers mention Monthly Cyclopædia

PEPPER WHISKY

ESTABLISHED 1780

THE GENUINE HAS THIS COUPON



PULL OFF ONLY When Bottle is Empty

EXTRA CAUTION
Anyone refilling this bottle without destroying the labels is criminally liable, and on detection will be prosecuted to the full extent of the law.

DISTILLERY BOTTLING
OLD PEPPER WHISKY
ESTABLISHED 1780
THE OLDEST & BEST BRAND OF WHISKY MADE IN KENTUCKY

Every Genuine BOTTLE HAS OUR TRADE MARK AND FURNISHES SIGNATURE BURNED ON THE HEAD

PEPPER DISTILLERY
HAND MADE
SOUR MASH
ESTABLISHED 1780
JAS. E. PEPPER & CO.
DISTILLERS,
LEXINGTON, KY.

Consumers SHOULD SURELY themselves that this WHISKY IS BOTTLED BY

Jas. E. Pepper & Co.
LEXINGTON, KY.

READ COUPON ON BACK OF THIS LABEL

FROM
HONEST DEALERS
OR THE
DISTILLERY DIRECT
AND THUS
AVOID IMITATIONS
AND REFILLED
BOTTLES

Genuine distillery bottling of Old Pepper Whisky, distilled only by Jas. E. Pepper & Co., Lexington, Ky., under the same formula for more than 100 years, is guaranteed absolutely the purest and best in the world. Sample case, \$15.00, sent on trial, which, if not perfectly satisfactory, can be returned and money will be refunded.

COPYRIGHT 1897 BY
THE STROBRIDGE
LITHO CO.
CINCINNATI & NEW YORK

Read and save coupons on genuine Distillery Bottling of Old Pepper Whisky and see who gets the \$5500 in 1898, in addition to \$1.00 per dozen.

Been giving satisfaction for 118 years---need we say more ?

In writing to advertisers mention Monthly Cyclopædia

WAMPOLE'S LIQUID WHEAT.

Each teaspoonful represents **one ounce** of all parts of the **whole wheat grain**, partly pre-digested and concentrated *in vacuo*; has sufficient **diastase** to digest five times its weight of dry starch, and contains one-eighth grain U. S. P. Pepsin.

Liquid Wheat contains over **seven per cent.** of the Nutrient Organic Hypophosphites, obtained solely from wheat, principally from the germ and cortical portions. Contains all the elements of nutrition, and is an efficient aid to the digestion of both **starchy** and **nitrogenous** foods. Valuable as a supplementary diet for growing children, insuring sufficient **tooth nutrition**, allaying nervous irritability, and inducing **plumpness**.


SAMPLES AND LITERATURE CHEERFULLY FURNISHED UPON APPLICATION.

Prepared by

HENRY K. WAMPOLE & CO.,

Manufacturing Pharmacists, Philadelphia, Pa.

"DO NOT STAMMER."



EDWIN S. JOHNSTON,
Principal and Founder

MR. E. BROWN.

DEAR SIR:—Answering your letter, just received: I am acquainted with E. S. Johnston, who used to be a frightful stammerer. He cured himself, and I have seen a great many others whom he has cured. I have confidence in his treatment.

Yours truly,

JOHN WANAMAKER.

Office of the Postmaster-General,
Washington, D. C., December 29, 1892.

CURED NINE YEARS—THE CASE OF WILLIS HART.

In reply to an inquiry letter, Mrs. Julia Hart, Oberlin, Ohio, writes: "In regard to Mr. E. S. Johnston's Institute, I can truly say that when I was there with my son, nine years ago, it was all it was recommended to be. I consider him a thoroughly reliable person. His dealings with me were gentlemanly, honest, and satisfactory. We shall never cease to be grateful to him for what he has done for our son. A lady said a few days ago: 'If I had not known Willis before his cure, it would be hard to believe that he ever stammered, and it would be hard for me to believe that he could ever have been cured.'"

Send for 60-page book to the PHILADELPHIA INSTITUTE, 1033 Spring Garden St., Philadelphia, Pa. Established 1884. EDWIN S. JOHNSTON, Principal and Founder (who cured himself after stammering 40 years).

A MOST INTERESTING NEW BOOK.

AN EPITOME OF THE HISTORY OF MEDICINE

From the Earliest Times to the Present.

By **ROSWELL PARK, A.M., M.D.,**

Professor of Surgery and Clinical Surgery in the Medical Department of the University of Buffalo,
Buffalo, New York, etc., etc.

Handsomely Illustrated with Half-tone Reproductions of Portraits of the Eminent
Physicians and Surgeons of all time and other Engravings.

Royal Octavo. 350 Pages. Extra Cloth. \$2.00 net. Now Ready.

THE F. A. DAVIS CO., Publishers, 1914-16 Cherry St., Philadelphia.

In writing to advertisers mention Monthly Cyclopædia

JUST PUBLISHED.

NEW EDITION OF PRINCIPLES OF SURGERY.

By N. SENN, M.D., Ph.D., LL.D.,

Professor of Practice of Surgery and Clinical Surgery in Rush Medical College, Chicago, Ill.; Professor of Surgery in the Chicago Polyclinic; Attending Surgeon to the Presbyterian Hospital; Surgeon-in-Chief to St. Joseph's Hospital, Chicago, Ill., etc.

Illustrated with **COLORED PLATES** and Numerous **WOOD-ENGRAVINGS**.

Second Edition. Thoroughly Revised and Enlarged. Royal Octavo. Nearly 700 Pages.

Net Prices: Extra Cloth, \$4.50; Sheep, \$5.50; Half-Russia, \$5.50.

These prices do not include Custom duties to purchasers in Canada.

THE POINTS ABOUT THIS WORK TO BE REMEMBERED ARE:

It is a New Edition.
Thoroughly Revised.
Considerably Enlarged.
With Sixty - nine Additional
Illustrations (a total of 178)
and Five Colored Plates.
Clear Type.
Excellent Paper.
Careful Presswork.
Substantial Bindings.
The Cost to the Physician and
Student is Most Moderate
and Reasonable.
It is a Scientific Work.

A Scholarly—that is, An Accu-
rately-Written—Work.
A Text-Book in Numerous Col-
leges.
Warmly Commended by Teach-
ers of Surgery Everywhere.
The Criticisms of the Press are
Universally Commendatory
and Full of Enthusiastic
Praise of the Whole Work.
Fully Abreast of the Times.
An Indispensable Addition to
the Working or Reference
Library of Every Live, Earn-
est Physician or Surgeon.

The subject-matter in this, the most important of the many able works from Dr. Senn's pen, includes: Regeneration. Regeneration of Different Tissues. Inflammation. Pathogenic Bacteria. Necrosis. Suppuration. Suppurative Osteomyelitis. Suppuration in Large Cavities. Abscess of Internal Organs. Ulcers and Fistula. Septicæmia. Pyæmia. Erysipelas. Tetanus. Hydrophobia. Surgical Tuberculosis. Tuberculosis of Lymphatic Glands, Peritoneum, Bones and Joints, Tendon-Sheaths, etc. Actinomycosis Hominis. Anthrax. Glanders, etc. A great deal of very valuable new material on the above and other subjects has been incorporated in this new edition, greatly enhancing its practical value.

THE F. A. DAVIS CO., Medical Publishers, 1914-16 Cherry St., Phila.

BRANCH OFFICES: { 117 W. Forty-Second St., New York City, U. S. A.
9 Lakeside Building, 214-220 S. Clark St., Chicago, Ill., U. S. A.

Salesmen Wanted.

Years ago the country merchant went to the city once or twice a year to buy his goods. Now a traveler calls on him every week, and he buys what he wants from the latest samples.

The traveler has come to stay.

Years ago the country doctor received catalogues and ordered therefrom, when money was so plentiful that he did not know what else to do with it. Consequently he did not order very much, and good libraries were few and far between.

Following the example of the dry-goods man, the medical publisher began sending out travelers about twenty-five years ago. Is the universally admitted fact, that medicine has made greater progress in that time than during a century before, a coincidence only?

However this may be, the fact is that the progressive country doctor, a hundred miles from any city, now keeps better informed than the city doctor did then.

He has become accustomed to having the latest ideas in medicine presented to him by the wide-awake traveler. He welcomes him as a source of reliable information on the best publications.

While he does not always buy, he has, like the dry-goods man, become accustomed to consider this method of buying the most satisfactory one, because he can examine his purchase thoroughly before he pays his money for it.

In this is the traveler of great use to the physician.

We claim to have the largest force of *expert* travelers in the business. We want a few more. Physicians who would prefer to hustle days rather than nights, can hear about it by addressing, with particulars,

THE SUBSCRIPTION DEPARTMENT,

THE F. A. DAVIS COMPANY,

1914-16 Cherry St., Philadelphia, Pa.

In writing to advertisers mention Monthly Cyclopædia

LIST OF PUBLICATIONS

OF

THE F. A. DAVIS COMPANY,

PHILADELPHIA: 1914-16 Cherry Street.

NEW YORK: 117 West Forty-Second Street.

CHICAGO: 9 Lakeside Building, 214-220 South Clark Street.

Age of the Domestic Animals. (Huidekoper)	\$ 1.00	Latin, Elements of. For Students of Medicine and Pharmacy. (Crothers and Bice)	\$ 1.25
Anatomy of the Central Nervous System of Man, and of Vertebrates in General. (Edinger)	2.50	Life-Insurance, Record-Book of Medical Examinations for. (Keating)	.50
Anatomy, Text-Book on Practical. (Boenning)	1.50	Lungs, Heart, and Kidneys, Diseases of the. (Davis)	1.25
Anatomy, Synopsis of Human. (Young)	1.00	Materia Medica, Pharmacy, and Therapeutics, Hand-Book of. (Bowen)	1.00
Annual and Analytical Cyclopædia of Practical Medicine, Sajous's . . . per vol.	5.00	Materia Medica and Therapeutics. (Shoemaker)	5.00
Annual of the Universal Medical Sciences: the Standard Medical Annual of the World	15.00	Medical Bulletin per year,	1.00
Auto-Intoxication in Disease. (Bouchard)	1.75	Medical Bulletin Visiting-List, or Physicians' Call-Record	1.25
Autoscopy of the Larynx and Trachea. (Kirstein)	.75	Medical Electro-Physics and Galvanism, International Text-Book of. (Bigelow, Editor)	2.50
Bacteriological Diagnosis. (Eisenberg)	1.00	Medical Formulary, International Pocket. (Witherstine)	2.00
Book on the Physician Himself and Things that Concern his Reputation and Success. (Cathell)	2.00	Medical Symbolism. (Sozinsky)	.75
Chart, Improved Clinical. (Bashore)	.50	Nervous Diseases, Lectures on. (Ranney)	3.75
Chinese: Their Present and Future. Medical, Political, and Social. (Coltman)	1.25	Nervo-Vascular System, Three Charts of the. (Price and Eagleton)	.50
Circumcision, History of. (Remondino)	1.25	Neuroses of the Genito-Urinary System in the Male. (Ultzmann)	1.00
Chapter on Cholera for Lay Readers. (Vought)	.50	Nose and Throat, Diseases of the. (Ivins)	4.00
Color-Vision and Color-Blindness. (Jennings)	1.00	Nose and Throat, Diseases of the. (Sajous)	3.00
Consumption. (Davis)	.75	Obstetrics, Practical. (Grandin and Jarmin)	4.00
Daughter, The: Her Health, Education, and Wedlock. (Capp)	.50	Obstetric Synopsis. (Stewart)	.75
Diabetes. (Purdy)	1.25	Oxygen and Other Gases in Medicine and Surgery. (Demarquay)	1.00
Ear, Nose, and Throat. Diseases of the. (Bishop)	4.00	Physicians' Interpreter. In English, French, German, and Italian	1.00
Eclampsia, Hand-Book of. (Michener)	.40	Physician's Wife, The. (Firebaugh)	1.25
Electro-Therapeutics, International System of. (Bigelow, Chief Editor)	6.00	Physicians' All-Requisite Time- and Labor-Saving Account Books	5.00
Electricity in Medicine and Surgery. (Liebig and Rohé)	2.00	Physiology of the Domestic Animals. (Smith)	3.00
Embryology, A Syllabus of Lectures on Human. (Manton)	1.25	Plain Talks on Avoided Subjects. (Guernsey)	1.00
Epilepsy. (Hare)	1.25	Primer of Psychology and Mental Disease. (Burr)	1.00
Eye-Strain in Health and Disease. (Ranney)	2.00	Psychopathia Sexualis. (Krafft-Ebing)	3.00
Eye, Injuries to the. (Baudry)	1.25	Rectum, Anus, and Contiguous Textures, Diseases of the. (Gant)	3.50
Face and Form Reading, The Encyclopedia of. (Stanton)	4.00	Skin, A Practical Text-Book on Diseases of the. (Rohé and Lord)	1.25
Fevers, Hand-Book of. (Eshner)	2.00	Southern California: Its Climate, Physical, and Meteorological Conditions. (Remondino)	.40
Fever. (Hare)	1.25	Spinal Concussion. (Clevenger)	1.50
Gynecology and Electro-Therapeutics, Conservative. (Massey)	3.50	Suggestive Therapeutics in Psychopathia Sexualis. (Schrenck-Notzing)	2.50
Gynecology, Lessons in. (Goodell)	2.50	Surgery, Principles of. (Senn)	4.50
Gynecology, Practical. (Montgomery) Prep'g.		Swedish Movement and Massage. (Nissen)	1.00
Hay Fever (Sajous)	.60	Symptom Register and Case Record. (Straub)	.75
Heredity, Health, and Personal Beauty. (Shoemaker)	2.50	Syphilis To-day and Among the Ancients. (Buret)	1.50
History of the Life of D. Hayes Agnew, M.D., LL. D. (Adams)	1.50	Tuberculosis of the Bones and Joints. (Senn)	4.00
History of Medicine, An Epitome of the. (Park)	2.00	Uranalysis and Urinary Diagnosis. (Purdy)	2.50
Hygiene, Text-Book of. (Rohé)	3.00		
Hygiene, Outlines of Rural. (Bashore)	.75		
Insanity. (Krafft-Ebing). In preparation.			

UNLESS OTHERWISE MENTIONED, BINDINGS ARE CLOTH AND PRICES ARE STRICTLY NET.

Complete Descriptive Catalogue Mailed on Request.

In writing to advertisers mention Monthly Cyclopædia

JUST PUBLISHED.

Conservative Gynecology and Electro-Therapeutics.

**A PRACTICAL TREATISE ON THE DISEASES OF WOMEN AND THEIR
TREATMENT BY ELECTRICITY.**

Third Edition. Revised, Rewritten, and Greatly Enlarged.

By G. BETTON MASSEY, M.D.,

Physician to the Gynecologic Department of Howard Hospital and late Electro-Therapeutist to the Infirmary for Nervous Diseases; Fellow and Ex-President of the American Electro-Therapeutic Association; Member of the Société Française d'Electrothérapie, American Medical Association, etc., etc.

Appropriately Illustrated with twelve (12) full-page handsome Chromo-Lithographic Plates, in twelve colors, of Drawings and Paintings made from nature specially for this work by a competent artist and with twelve (12) beautifully-printed full-page Half-tone Plates of Photographs taken from nature, as well as 140 Photo- and Wood-engravings in the text.

**COMPLETE IN ONE ROYAL OCTAVO VOLUME. BOUND IN EXTRA CLOTH, BEVELED EDGES. 400 PAGES.
PRICE, \$3.50 NET, PUBLISHERS PAYING DELIVERY CHARGES.**

This work is not only a revision and expansion of the author's two preceding editions of "Electricity in the Diseases of Women," with a full consideration of the latest advances in the application of electricity in gynecology, but it has been recast in the form of a complete treatise on Conservative Gynecology from the Stand-point of the General Practitioner. Recognizing the fact that most modern treatises on gynecology are written from the point of view of the operative surgeon, the author has endeavored to produce a work which will assist the physician in the management of these cases without recourse to the knife, whenever this is practicable, and deals fully with the indications for the employment of electricity after the methods of Apostoli and others.

The work is, therefore, an appeal for a more conservative treatment of the diseases of women, and presents ample data in justification of the author's well-known position on this point, followed by detailed directions for the application of the methods described, together with a complete introduction to medical electricity. In a chapter on malignant affections a full description is given of the method discovered by the author whereby mercury and other chemically nascent lethal substances are disseminated throughout a cancer by cataphoresis for the interstitial destruction of malignant cells.

THE F. A. DAVIS COMPANY, Medical Publishers.

PHILADELPHIA: 1914-16 Cherry Street.

NEW YORK: 117 W. Forty-second Street.

CHICAGO: 9 Lakeside Building, 214-220 S. Clark Street.

In writing to advertisers mention Monthly Cyclopædia

MEDICAL STUDENTS

should bear in mind that while a knowledge of anatomy and physiology may be essential to their success as practitioners,

A KNOWLEDGE OF BUSINESS IS

still more so, as no physician can expect to do himself justice, professionally, who is

HARASSED BY FINANCIAL DIFFICULTIES.

To know anything of business, a man must actually engage in business.

THE PHYSICIAN'S ALL-REQUISITE ACCOUNT-BOOK

is a good seller, and the commission for taking orders is liberal. Money can be made right at home, in every city or large town. Customers can be secured and a regular trade established.

Write for particulars.

THE F. A. DAVIS COMPANY,

1914 and 1916 Cherry Street,

Philadelphia, Pa.

SOUND ADVICE.

"I have not become acquainted with Dr. Cathell through the medical press; I do not know whether he has discovered a new bacterium, or a novel method of emptying the female pelvis, or an antitubercular serum; but I do believe in this book he will live in the grateful hearts of generation after generation of physicians who will owe much of their success to him. He has seen much, and his experience is here crystallized into sound advice. There is help in it for the physician who finds, after a short time, that 'something is wrong' with his career. Here he can find what is the matter and the remedy for it. The senior student should read it before he even thinks of a location; he should read it often during his early professional life, for from it he will gain much that he would otherwise obtain only from sometimes bitter experience."

G. H. S. in "The Colorado Medical Journal,"
Denver, Colorado.

THE FOREGOING REMARKS REFER TO THE BOOK ON THE PHYSICIAN HIMSELF,

And Things that Concern his Reputation
and Success.

— BY —

D. W. CATHELL, M.D.

Tenth Edition. Author's last revision. Royal Octavo. About 350
pages. Extra Cloth. Net, \$2.00.

THE F. A. DAVIS CO., Publishers.

PHILADELPHIA: 1914 and 1916 Cherry Street.

NEW YORK: 117 W. Forty-Second Street.

CHICAGO: 9 Lakeside Building, 214-220 S. Clark Street.

In writing to advertisers mention Monthly Cyclopædia

STANDARD MODERN MEDICAL TEXT-BOOKS

ISSUED BY

THE F. A. DAVIS COMPANY, Medical Publishers,

PHILADELPHIA, PA.

Bindings in Cloth, Except when Otherwise Stated.

ALL THE FOLLOWING PRICES ARE STRICTLY NET. CARRIAGE CHARGES PREPAID BY THE PUBLISHERS.

- *BASHORE—*Outlines of Rural Hygiene.* 75 cents.
- *BISHOP—*Diseases of the Ear, Nose, and Throat.* Second Edition. Cloth, \$4.00; Sheep or Half-Russia, \$5.00.
- BOUCHARD—OLIVER—*Auto-Intoxication in Disease. A Treatise on Intestinal and Urinary Pathology.* \$1.75.
- BURR—*Primer of Psychology and Mental Disease.* Second Edition, \$1 00.
- CATHELL—*Book on the Physician Himself.* Tenth Edition, \$2.00.
- CROTHERS AND BICE—*Elements of Latin for Students in Medicine and Pharmacy.* \$1.25.
- *EDINGER—HALL—*Structure of the Central Nervous System.* New, Enlarged, American Edition, \$2.50
- EISENBERG—PIERCE—*Bacteriological Diagnosis.* Cloth or Oil-cloth, \$1.00.
- *GANT—*Diseases of the Rectum, Anus, and Contiguous Textures.* Cloth, \$3.50; Half-Russia, \$4.50
- *GRANDIN AND JARMAN—*Practical Obstetrics.* Second Edition. Cloth, \$4.00; Sheep, \$4 75.
- **International System of Electro-Therapeutics.* Edited by H. R. BIGELOW, M.D. Cloth, \$6.00; Sheep \$7.00; Half-Russia, \$7.50.
- **International Text-Book of Medical Electro-Physics and Galvanism.* \$2.50.
- *IVINS—*Diseases of the Nose and Throat.* Cloth, \$4.00. Sheep or Half-Russia, \$5.00.
- *MANTON—*Syllabus of Lectures on Human Embryology.* Second Edition, \$1.25.
- *MASSEY—*Conservative Gynecology and Electro-Therapeutics.* Third Edition, \$3.50.
- *MONTGOMERY—*A Text-book of Practical Gynecology.* IN PREPARATION.
- *PARK—*An Epitome of the History of Medicine.* \$2.00.
- PRICE AND EAGLETON—*Three Charts of the Nervo-Vascular System, Nerves, Arteries, and Veins.* Unmounted, on extra durable paper. Per Set, 50 cents.
- *PURDY—*Practical Urinalysis and Urinary Diagnosis.* Fourth Edition, \$2.50.
- *ROHÉ—*Text-book of Hygiene.* Third Edition, \$3.00.
- *ROHÉ AND LORD—*Practical Manual of Skin Diseases.* \$1.25.
- *SENN—*Principles of Surgery.* Second Edition. Cloth, \$4.50; Sheep or Half-Russia, \$5.50.
- SHOEMAKER—*Materia Medica and Therapeutics.* Fourth Edition. Cloth, \$5.00; Sheep, \$5.75.
- *SMITH—*Physiology of the Domestic Animals.* Cloth, \$3.00; Sheep, \$4.00.
- *YOUNG—*Synopsis of Human Anatomy.* \$1.00.

* ALL BOOKS DESIGNATED BY AN ASTERISK (*) ARE APPROPRIATELY AND LIBERALLY ILLUSTRATED

DETAILED DESCRIPTIVE CIRCULARS OF ANY OF THESE BOOKS FORWARDED PROMPTLY ON REQUEST. CORRESPONDENCE WITH COLLEGE PROFESSORS AND TEACHERS CONCERNING ANY OF THE ABOVE BOOKS SPECIALLY SOLICITED.

THE F. A. DAVIS COMPANY, Publishers,

1914 and 1916 Cherry St., Philadelphia.

In writing to advertisers mention Monthly Cyclopædia

IMPORTANT ANNOUNCEMENT

Concerning DR. WINFIELD S. HALL'S Translation of the Latest German Edition of Edinger's "Structure of the Central Nervous System."

IN PRESS. WILL BE READY SOON.

THE ANATOMY OF THE Central Nervous System of Man and of Vertebrates in General.

By Prof. LUDWIG EDINGER, M.D.,
Frankfort-on-the-Main, Germany.

Translated from the FIFTH GERMAN EDITION,

By WINFIELD S. HALL, Ph.D., M.D.,
Professor of Physiology in the Northwestern University Medical School, Chicago.

ASSISTED BY

PHILO LEON HOLLAND, M.D., Instructor in Clinical Neurology, and
EDWARD P. CARLTON, B.S., Demonstrator of Histological Neurology in the Northwestern
University Medical School, Chicago.

Professors and Teachers of Anatomy of the Nervous System, Histology, etc., in Medical Colleges throughout the English-speaking world will be gratified to learn that the above book will be ready in time for the college sessions of '98 and '99.

This translation is a thoroughly complete one from the latest German edition, admirably executed, and done into clear, correct, and readable English with the authority and indorsement of the famous author. It contains upward of 250 engravings in the text and comprises over 400 Royal Octavo pages. It will be handsomely printed and bound in extra cloth, and sold at the exceedingly moderate price of \$2 50 net.

That this is one of the most important, interesting, and useful books which can be placed in the hands of a medical student during his attendance at college will be quickly recognized by every Teacher of Anatomy, Histology, and Neurology.

Further information promptly furnished by the publishers.

THE F. A. DAVIS COMPANY, Publishers.

PHILADELPHIA: 1914 and 1916 Cherry St.

NEW YORK: 117 W. Forty-Second St.

CHICAGO: 9 Lakeside Building, 214-220 S. Clark St.

In writing to advertisers mention Monthly Cyclopædia

SYR. HYPOPHOS. CO., FELLOWS

Contains the Essential Elements of the Animal Organization—Potash and Lime;

The Oxidising Agents—Iron and Manganese;

The Tonics—Quinine and Strchnine;

And the Vitalizing Constituent—Phosphorus; the whole combined in the form of a Syrup with a Slightly Alkaline Reaction.

It Differs in its Effects from all Analogous Preparations; and it possesses the important properties of being pleasant to the taste, easily borne by the stomach, and harmless under prolonged use.

It has Gained a Wide Reputation, particularly in the treatment of Pulmonary Tuberculosis, Chronic Bronchitis, and other affections of the respiratory organs. It has also been employed with much success in various nervous and debilitating diseases.

Its Curative Power is largely attributable to its stimulant, tonic, and nutritive properties, by means of which the energy of the system is recruited.

Its Action is Prompt; it stimulates the appetite and the digestion, it promotes assimilation, and it enters directly into the circulation with the food products.

The prescribed dose produces a feeling of buoyancy, and removes depression and melancholy; *hence the preparation is of great value in the treatment of mental and nervous affections.* From the fact, also, that it exerts a double tonic influence, and induces a healthy flow of the secretions, its use is indicated in a wide range of diseases.

Medical Letters may be addressed to **Mr. FELLOWS, 48 Vesey Street, New York.**



Open All the Year. 2,500 FEET ELEVATION.
ON CHESAPEAKE & OHIO RAILWAY.

THE NEW HOMESTEAD,

a grand new hotel completed last year, equipped and furnished throughout in accordance with the latest designs and most modern improvements. The Virginia, the Annex, and ten new cottages clustered near the Homestead. A handsome new Casino adds greatly to indoor enjoyment.

*Golf Links and Club House, Tennis Courts, Billiards, Pool,
Riding School and Fine Livery.*

MAGNIFICENT MOUNTAIN SURROUNDINGS.

FINEST BATH-HOUSE IN AMERICA.

MOST CURATIVE BATHS KNOWN

for rheumatism, gout, and nervous troubles. All baths from flowing streams of natural hot water.

Located nearly midway between Washington, Old Point Comfort, and Cincinnati or Louisville. Hot Springs is an exceptionally convenient and delightful stopping-place for west-bound travelers.

Special Pullman Compartment Car leaves Pennsylvania Station, New York, 4.50 P.M., Tuesday, Thursday, and Saturday, and arrives at Hot Springs via Washington, 7.25 A.M. Dining car.

Excursion tickets, baggage checks, and Pullman Reservations at Chesapeake & Ohio Railway Offices, 362 and 1323, Broadway, New York, also offices Pennsylvania Railroad and connecting lines.


For illustrated pamphlets and further particulars, address W. H. GREGOR, General Agent Passenger Department Chesapeake and Ohio Railway, Washington, D. C., or

FRED STERRY, Manager, Hot Springs, Bath Co., Va.

In writing to advertisers mention Monthly Cyclopædia

Oak Grove Hospital


For Nervous and Mental Diseases.



Equipment for hydrotherapeutic and electrical treatment complete and modern—Static, Galvanic, and Faradic Apparatus, Electric Bath, Turkish and Russian Baths, and Massage. Noyes Amusement Hall in connection with the Hospital, containing gymnasium, billiard-room, bowling-alley, hall for assemblies, sitting-room, lounging-room, and library, furnishes ideal facilities for in-door sports and diversion. Carriages for the daily use of patients free. For terms, address

Grounds comprise sixty acres of stately oaks and are picturesque and secluded. Buildings roomy, homelike, free from institutional features, and erected with especial reference to the care of nervous and mental invalids.

Interiors Bright and Cheerful, Luxurious Furnishings, Superior Appointments, Skilled Attendance, First-class Cuisine.



Dr. C. B. BURR, Medical Director,
FLINT, MICH.

BLOOD VITALITY

The Essence of Life is the blood.

The Vital Element of the blood is Hæmoglobin.

Without a normal percentage of this elementary principle the tissues are insufficiently oxygenated and poorly nourished.

With a proper proportion the vital functions are quickened and the entire system thoroughly fortified.

Pepto-Mangan ("Gude")

is a hæmoglobin producer, because of the quickly assimilable organic Iron and Manganese which it contributes to the devitalized circulatory fluid existing in cases of

Anæmia, Chlorosis, Amenorrhœa, Dysmenorrhœa,
Bright's Disease, Chorea, etc.

Prescribe PEPTO-MANGAN "GUDE" in original bottles containing $\frac{3}{4}$ xi.

IT'S NEVER SOLD IN BULK.

M. J. BREITENBACH COMPANY,

Sole Agents for United States and Canada,

LABORATORY:

LEIPZIG, GERMANY.

56-58 WARREN ST., NEW YORK.



MARCHAND'S EYE BALSAM

(C. P. Vegetable Glycerine combined with Ozone)

IS THE MOST POWERFUL AND AT THE SAME TIME HARMLESS
HEALING AGENT KNOWN.

HYDROZONE (30 volumes preserved aqueous solution of H_2O_2)

IS THE MOST POWERFUL ANTISEPTIC AND PUS DESTROYER.
HARMLESS STIMULANT TO HEALTHY GRANULATIONS.

Cure quickly Suppurative and Inflammatory Diseases of the Eye:

**Catarrhal Conjunctivitis or Ophthalmia,
Purulent Conjunctivitis, — Ophthalmia in Children,
Inflamed and Granular Eye Lids, Etc.**

Send for free 240-page book "Treatment of Diseases caused by Germs," containing reprints of 120 scientific articles by leading contributors to medical literature.

Physicians remitting 50 cents will receive one complimentary sample of each, "Hydrozone" and "Eye Balsam" by express, charges prepaid.

Marchand's Eye Balsam is put up only in one size bottle. Package sealed with my signature.

Hydrozone is put up only in extra small, small, medium, and large size bottles, bearing a red label, white letters, gold and blue border with my signature.

Glycozone is put up only in 4-oz., 8-oz. and 16-oz. bottles, bearing a yellow label, white and black letters, red and blue border with my signature.

PREPARED ONLY BY

Charles Marchand

Chemist and Graduate of the "Ecole Centrale
des Arts et Manufactures de Paris" (France).

Charles Marchand,

28 Prince St., New York.

Sold by leading Druggists.

Avoid Imitations.

☞ Mention this Publication.

When Ordering Iron as a Tonic

for child, woman, or man, to achieve increase of
hemoglobin and red blood corpuscles, in cases of

Anemia, Chlorosis, Malnutrition,

or any debilitated condition, after fevers and dur-
ing convalescence from any affection, : : : : :

Specify Ferratin

It is the only organic iron compound the therapeutic application
of which is based on physiological tests (record of which is con-
tained in all Annuals and Text-books issued since 1894). : : : :

FERRATIN TABLETS, $7\frac{1}{2}$ grains
each. Palatable and most con-
venient. Sold by all Druggists...
Order an original box of 50 tablets.

Literature Mailed on Request.

C. F. BOEHRINGER & SOEHNE,
7 CEDAR STREET, NEW YORK.